

Anti-p53R2 Antibody

Catalog # ABO11361

Product Information

Application	WB
Primary Accession	Q7LG56
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Ribonucleoside-diphosphate reductase subunit M2 B(RRM2B) detection. Tested with WB in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	50484
Other Names	Ribonucleoside-diphosphate reductase subunit M2 B, 1.17.4.1, TP53-inducible ribonucleotide reductase M2 B, p53-inducible ribonucleotide reductase small subunit 2-like protein, p53R2, RRM2B, P53R2
Calculated MW	40737
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse
Subcellular Localization	Cytoplasm. Nucleus. Translocates from cytoplasm to nucleus in response to DNA damage.
Tissue Specificity	Widely expressed at a high level in skeletal muscle and at a weak level in thymus. Expressed in epithelial dysplasias and squamous cell carcinoma. .
Source	Eukaryota
Protein Name	Ribonucleoside-diphosphate reductase subunit M2 B
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human p53R2(318-332aa EGKTNFFEKRVSEYQ), identical to the related rat and mouse sequences.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins

Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the ribonucleoside diphosphate reductase small chain family.


Protein Information

Name	RRM2B
Synonyms	P53R2
Function	Plays a pivotal role in cell survival by repairing damaged DNA in a p53/TP53-dependent manner. Supplies deoxyribonucleotides for DNA repair in cells arrested at G1 or G2. Contains an iron-tyrosyl free radical center required for catalysis. Forms an active ribonucleotide reductase (RNR) complex with RRM1 which is expressed both in resting and proliferating cells in response to DNA damage.
Cellular Location	Cytoplasm. Nucleus. Note=Translocates from cytoplasm to nucleus in response to DNA damage
Tissue Location	Widely expressed at a high level in skeletal muscle and at a weak level in thymus. Expressed in epithelial dysplasias and squamous cell carcinoma.

Background

RRM2B(Ribonucleotide Reductase, M2 B), also known as P53R2, is an enzyme that in humans is encoded by the RRM2B gene. Tanaka et al.(2000) mapped the p53R2 gene to chromosome 8q23.1 by fluorescence in situ hybridization. Tanaka et al.(2000) found that expression of p53R2, but not R2, was induced by ultraviolet and gamma-irradiation and adriamycin treatment in a wildtype p53-dependent manner. Induction of p53R2 in p53-deficient cells caused G2/M arrest and protected cells from death in response to adriamycin. Inhibition of endogenous p53R2 expression in cells that had an intact p53-dependent DNA damage checkpoint reduced ribonucleotide reductase activity, DNA repair, and cell survival after exposure to various genotoxins.

Images



Anti-p53R2 antibody, ABO11361, Western blotting
Lane 1: Rat Thymus Tissue Lysate
Lane 2: MCF-7 Cell Lysate
Lane 3: A431 Cell Lysate
Lane 4: HELA Cell Lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.