10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-PMP70 Antibody

Catalog # ABO11398

Product Information

Application	WB, IHC-P
Primary Accession	P28288
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for ATP-binding cassette sub-family D member 3(ABCD3) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	5825
Other Names	ATP-binding cassette sub-family D member 3, 70 kDa peroxisomal membrane protein, PMP70, ABCD3, PMP70, PXMP1
Calculated MW	75476
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 □g/ml, Human, Rat, Mouse, By Heat Western blot, 0.1-0.5 □g/ml, Rat, Human, Mouse
Subcellular Localization	Peroxisome membrane ; Multi-pass membrane protein .
Source	Eukaryota
Protein Name	ATP-binding cassette sub-family D member 3
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human PMP70(646-659aa EFKQITEDTVEFGS), different from the related mouse and rat sequences by one amino acid.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20˚C for one year. After r˚Constitution, at 4˚C for one month. It˚Can also be aliquotted and stored frozen at -20˚C for a longer time.Avoid repeated

	freezing and thawing.
Sequence Similarities	Belongs to the ABC transporter superfamily. ABCD family. Peroxisomal fatty acyl CoA transporter (TC 3.A.1.203) subfamily.

Protein Information

Name	ABCD3 {ECO:0000303|PubMed:39223112, ECO:0000312|HGNC:HGNC:67}
Function	ATP-dependent transporter of the ATP-binding cassette (ABC) family that catalyzes the transport of long- and very-long-chain fatty acyl-CoAs and probably bile acyl-CoAs from the cytosol to the peroxisome matrix for beta-oxydation. Appears to be specifically involved in the peroxisomal transport of branched-chain fatty acyl-CoAs (BCFA-CoAs) and dicarboxylic acyl CoAs (DCA-CoAs) (PubMed:11248239, PubMed:23671276, PubMed:24333844, PubMed:25168382, PubMed:29397936, PubMed:39223112, PubMed:41428872). Has fatty acyl-CoA thioesterase and ATPase activities. Probably hydrolyzes fatty acyl-CoAs into free fatty acids prior to their ATP-dependent transport into peroxisomes.
Cellular Location	Peroxisome membrane; Multi-pass membrane protein


Background

ATP-binding cassette sub-family D member 3 is a protein that in humans is encoded by the ABCD3 gene. The protein encoded by this gene is a member of the superfamily of ATP-binding cassette(ABC) transporters. The gene was assigned to human chromosome 1p21-p22 by in situ hybridization. This peroxisomal membrane protein likely plays an important role in peroxisome biogenesis. Mutations have been associated with some forms of Zellweger syndrome, a heterogeneous group of peroxisome assembly disorders.


Images



Anti-PMP70 antibody, ABO11398, Western blottingLane 1: Rat Lung Tissue LysateLane 2: Rat Ovary Tissue Lysate



Anti-PMP70 antibody, ABO11398, IHC(P)IHC(P): Human Lung Cancer Tissue



Anti-PMP70 antibody, ABO11398, IHC(P)IHC(P): Rat Brain Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.