

Anti-SMAD5 Antibody

Catalog # ABO11422

Product Information

Application	WB, IHC-P
Primary Accession	Q99717
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Mothers against decapentaplegic homolog 5(SMAD5) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	4090
Other Names	Mothers against decapentaplegic homolog 5, MAD homolog 5, Mothers against DPP homolog 5, JV5-1, SMAD family member 5, SMAD 5, Smad5, hSmad5, SMAD5, MADH5
Calculated MW	52258
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Rat, Mouse, By Heat Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse
Subcellular Localization	Cytoplasm. Nucleus. Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4.
Tissue Specificity	Ubiquitous.
Source	Eukaryota
Protein Name	Mothers against decapentaplegic homolog 5
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg Na ₃ .
Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human SMAD5(248-267aa IPQIMPSSSRDVQPVAYEE), identical to the related rat sequence, and different from the related mouse sequence by one amino acid.
Purification	Immunogen affinity purified.

Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the dwarfin/SMAD family.


Protein Information

Name	SMAD5 (HGNC:6771)
Synonyms	MADH5
Function	Transcriptional regulator that plays a role in various cellular processes including embryonic development, cell differentiation, angiogenesis and tissue homeostasis (PubMed: 12064918 , PubMed: 16516194). Upon BMP ligand binding to their receptors at the cell surface, is phosphorylated by activated type I BMP receptors (BMPRIIs) and associates with SMAD4 to form a heteromeric complex which translocates into the nucleus acting as transcription factor (PubMed: 9442019). In turn, the hetero-trimeric complex recognizes cis- regulatory elements containing Smad Binding Elements (SBEs) to modulate the outcome of the signaling network (PubMed: 33510867). Non-phosphorylated SMAD5 has a cytoplasmic role in energy metabolism regulation by promoting mitochondrial respiration and glycolysis in response to cytoplasmic pH changes (PubMed: 28675158). Mechanistically, interacts with hexokinase 1/HK1 and thereby accelerates glycolysis (PubMed: 28675158).
Cellular Location	Cytoplasm. Nucleus Mitochondrion. Note=Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4
Tissue Location	Ubiquitous.


Background

Mother against decapentaplegic homolog 5 also known as SMAD5 is a protein that in humans is encoded by the SMAD5 gene. It belongs to the SMAD family of proteins, which belong to the TGFbeta superfamily of modulators. The gene was assigned to human chromosome 5q31. Like many other TGFbeta family members SMAD5 is involved in cell signalling and modulates signals of bone morphogenetic proteins(BMP's). It may play a role in the pathway where TGFbeta is an inhibitor of hematopoietic progenitor cells.


Images




Anti-SMAD5 antibody, ABO11422, IHC(P)IHC(P): Rat Intestine Tissue



Anti-SMAD5 antibody, ABO11422, IHC(P)IHC(P): Human Intestinal Cancer Tissue



Anti-SMAD5 antibody, ABO11422, IHC(P)IHC(P): Human Lung Cancer Tissue



Anti-SMAD5 antibody, ABO11422, Western blottingAll lanes: Anti SMAD5 (ABO11422) at 0.5ug/mlLane 1: K562 Whole Cell Lysate at 40ugLane 2: JURKAT Whole Cell Lysate at 40ugLane 3: PC 1-2 Whole Cell Lysate at 40ugLane 4: HELA Whole Cell Lysate at 40ugLane 5: SMMC Whole Cell Lysate at 40ugPredicted bind size: 52 KDObserved bind size: 52KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.