

Anti-Wnt4 Antibody

Catalog # ABO11427

Product Information

Application	WB
Primary Accession	P56705
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Protein Wnt-4(WNT4) detection. Tested with WB in Human;Mouse;Rat.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	54361
Other Names	Protein Wnt-4, WNT4
Calculated MW	39052
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat
Subcellular Localization	Secreted, extracellular space, extracellular matrix.
Source	Eukaryota
Protein Name	Protein Wnt-4
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Wnt4(232-249aa HALKEKFDGATEVEPRRV), identical to the related mouse and rat sequences.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20° C for one year. After reconstitution, at 4° C for one month. It can also be aliquotted and stored frozen at -20° C for a longer time. Avoid repeated freezing and thawing.

Protein Information

Name	WNT4
Function	Ligand for members of the frizzled family of seven transmembrane receptors (Probable). Plays an important role in the embryonic development of the urogenital tract and the lung (PubMed: 15317892 , PubMed: 16959810 , PubMed: 18179883 , PubMed: 18182450). Required for normal mesenchyme to epithelium transition during embryonic kidney development. Required for the formation of early epithelial renal vesicles during kidney development (By similarity). Required for normal formation of the Mullerian duct in females, and normal levels of oocytes in the ovaries (PubMed: 15317892 , PubMed: 16959810 , PubMed: 18182450). Required for normal down-regulation of 3 beta-hydroxysteroid dehydrogenase in the ovary (PubMed: 15317892 , PubMed: 16959810 , PubMed: 18182450). Required for normal lung development and for normal patterning of tracheal cartilage rings (By similarity).
Cellular Location	Secreted, extracellular space, extracellular matrix

Background

Wingless-type MMIV integration site family, member4, is a secreted protein that in humans is encoded by the Wnt4 gene. By fluorescence in situ hybridization, WNT4 gene was mapped to chromosome 1p35, a location consistent with the results of radiation hybrid mapping. The WNT gene family consists of structurally related genes that encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and embryogenesis.

Images



Anti- WNT4 antibody, ABO11427, Western blottingAll lanes: Anti WNT4 (ABO11427) at 0.5ug/mlWB: MCF-7 Whole Cell Lysate at 40ugPredicted bind size: 37KDObserved bind size: 37KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.