

Anti-EAAT3 Antibody

Catalog # ABO11476

Product Information

Application	WB
Primary Accession	P51906
Host	Rabbit
Reactivity	Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Excitatory amino acid transporter 3(SLC1A1) detection. Tested with WB in Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	20510
Other Names	Excitatory amino acid transporter 3, Excitatory amino-acid carrier 1, Sodium-dependent glutamate/aspartate transporter 3, Solute carrier family 1 member 1, Slc1a1, Eaac1, Eaat3
Calculated MW	56694
Application Details	Western blot, 0.1-0.5 µg/ml, Mouse, Rat
Subcellular Localization	Cell membrane ; Multi-pass membrane protein. Apical cell membrane ; Multi-pass membrane protein. Membrane; Multi-pass membrane protein.
Tissue Specificity	Expressed in brain, lung, kidney and skeletal muscle.
Source	Eukaryota
Protein Name	Excitatory amino acid transporter 3
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of mouse EAAT3(10-27aa DWRRFLRNHWLLSTVAA), identical to the related rat sequence.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins

Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
----------------	---

Protein Information


Name	Slc1a1
Synonyms	Eaac1 {ECO:0000303 PubMed:9233792}, Eaat
Function	Sodium-dependent, high-affinity amino acid transporter that mediates the uptake of L-glutamate and also L-aspartate and D-aspartate (PubMed: 12119102 , PubMed: 18684713). Can also transport L-cysteine (PubMed: 30840898). Functions as a symporter that transports one amino acid molecule together with two or three Na(+) ions and one proton, in parallel with the counter-transport of one K(+) ion. Mediates Cl(-) flux that is not coupled to amino acid transport; this avoids the accumulation of negative charges due to aspartate and Na(+) symport (By similarity). Plays an important role in L-glutamate and L-aspartate reabsorption in renal tubuli (PubMed: 9233792). Plays a redundant role in the rapid removal of released glutamate from the synaptic cleft, which is essential for terminating the postsynaptic action of glutamate (PubMed: 9233792). Contributes to glutathione biosynthesis and protection against oxidative stress via its role in L-glutamate and L- cysteine transport (PubMed: 30840898). Negatively regulated by ARL6IP5 (PubMed: 12119102).
Cellular Location	Cell membrane; Multi-pass membrane protein {ECO:0000250 UniProtKB:P43003}. Apical cell membrane; Multi-pass membrane protein {ECO:0000250 UniProtKB:P43003}. Synapse, synaptosome. Early endosome membrane. Late endosome membrane. Recycling endosome membrane
Tissue Location	Detected on neurons in the brain cortex, dentate gyrus and hippocampus CA2 region (at protein level) (PubMed:30840898) Expressed in whole brain, brain cortex, hippocampus, cerebellum, lung, kidney, small intestine and skeletal muscle (PubMed:30840898, PubMed:7766664, PubMed:9233792). Expressed in the renal outer medulla, medullary ray and cortex (at protein level) (PubMed:26739563)

Background

Solute carrier family 1 member 1, also called SLC1A1, is a protein that in humans is encoded by the SLC1A1 gene. By Southern analysis of a panel of human/rodent somatic cell hybrids and by fluorescence in situ hybridization(FISH), this gene is mapped to 9p24.2. This gene encodes a member of the high-affinity glutamate transporters that play an essential role in transporting glutamate across plasma membranes. In brain, these transporters are crucial in terminating the postsynaptic action of the neurotransmitter glutamate, and in maintaining extracellular glutamate concentrations below neurotoxic levels. This transporter also transports aspartate, and mutations in this gene are though to cause dicarboxylicamino aciduria, also known as glutamate-aspartate transport defect.

Images

Anti-EAAT3 antibody, ABO11476, Western blottingLane 1:
Rat Liver Tissue LysateLane 2: Rat Heart Tissue
LysateLane 3: HEPA Cell Lysate



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.