

Anti-IGF2R Antibody

Catalog # ABO11492

Product Information

Application	WB, IHC-P, ICC
Primary Accession	P11717
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Cation-independent mannose-6-phosphate receptor(IGF2R) detection. Tested with WB, IHC-P, ICC in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	3482
Other Names	Cation-independent mannose-6-phosphate receptor, CI Man-6-P receptor, CI-MPR, M6PR, 300 kDa mannose 6-phosphate receptor, MPR 300, Insulin-like growth factor 2 receptor, Insulin-like growth factor II receptor, IGF-II receptor, M6P/IGF2 receptor, M6P/IGF2R, CD222, IGF2R, MPRI
Calculated MW	274375
Application Details	Immunocytochemistry , 0.5-1 µg/ml, Human, Mouse, Rat Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Rat, Mouse, By Heat Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat
Subcellular Localization	Lysosome membrane ; Single-pass type I membrane protein . Colocalized with DPP4 in internalized cytoplasmic vesicles adjacent to the cell surface.
Source	Eukaryota
Protein Name	Cation-independent mannose-6-phosphate receptor
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg Na ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Mannose 6 Phosphate Receptor(Cation independent)(2475-2491aa TKLVSFHDDSDDLLHI), different from the related rat and mouse sequences by one amino acid.
Purification	Immunogen affinity purified.

Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the MRL1/IGF2R family.


Protein Information

Name	IGF2R
Synonyms	MPRI
Function	Mediates the transport of phosphorylated lysosomal enzymes from the Golgi complex and the cell surface to lysosomes (PubMed: 18817523 , PubMed: 2963003). Lysosomal enzymes bearing phosphomannosyl residues bind specifically to mannose-6-phosphate receptors in the Golgi apparatus and the resulting receptor-ligand complex is transported to an acidic prelysosomal compartment where the low pH mediates the dissociation of the complex (PubMed: 18817523 , PubMed: 2963003). The receptor is then recycled back to the Golgi for another round of trafficking through its binding to the retromer (PubMed: 18817523). This receptor also binds IGF2 (PubMed: 18046459). Acts as a positive regulator of T-cell coactivation by binding DPP4 (PubMed: 10900005).
Cellular Location	Golgi apparatus membrane; Single-pass type I membrane protein. Endosome membrane; Single-pass type I membrane protein. Note=Mainly localized in the Golgi at steady state and not detectable in lysosome (PubMed:18817523) Colocalized with DPP4 in internalized cytoplasmic vesicles adjacent to the cell surface (PubMed:10900005).


Background

Insulin-like growth factor 2 receptor, also called IGF2R or I-MPR is a protein that in humans is encoded by the IGF2R gene. This gene is mapped to 6q25.3. This gene encodes a receptor for both insulin-like growth factor 2 and mannose 6-phosphate, although the binding sites for either are located on different segments of the receptor. This receptor functions in the intracellular trafficking of lysosomal enzymes, the activation of transforming growth factor beta, and the degradation of insulin-like growth factor 2. While the related mouse gene shows exclusive expression from the maternal allele, imprinting of the human gene appears to be polymorphic, with only a minority of individuals showing expression from the maternal allele.


Images



Anti-Mannose 6 Phosphate Receptor(Cation independent) antibody, ABO11492, IHC(P)IHC(P): Human Intestinal Cancer Tissue



Anti-Mannose 6 Phosphate Receptor(Cation independent) antibody, ABO11492, ICCICC: HCT116 Cell



Anti-Mannose 6 Phosphate Receptor(Cation independent) antibody, Western blotting
All lanes: Anti Mannose 6 Phosphate Receptor (Cation independent) (ABO11492) at 0.5ug/ml
Lane 1: HELA Whole Cell Lysate at 40ug
Lane 2: JURKAT Whole Cell Lysate at 40ug
Lane 3: 22RV1 Whole Cell Lysate at 40ug
Lane 4: 293T Whole Cell Lysate at 40ug
Predicted bind size: 274KD
Observed bind size: 274KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.