

Anti-Alpha 2A Adrenergic Receptor Antibody

Catalog # ABO11507

Product Information

Application	WB
Primary Accession	P08913
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Alpha-2A adrenergic receptor(ADRA2A) detection. Tested with WB in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	150
Other Names	Alpha-2A adrenergic receptor, Alpha-2 adrenergic receptor subtype C10, Alpha-2A adrenoreceptor, Alpha-2A adrenoceptor, Alpha-2AAR, ADRA2A, ADRA2R, ADRAR
Calculated MW	50647
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat
Subcellular Localization	Cell membrane ; Multi-pass membrane protein .
Source	Eukaryota
Protein Name	Alpha-2A adrenergic receptor
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human alpha 2a Adrenergic Receptor(213-228aa ILVYVRIYQIAKRRT), identical to the related mouse and rat sequences.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20° C for one year. After reconstitution, at 4° C for one month. It can also be aliquotted and stored frozen at -20° C for a longer time.Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the G-protein coupled receptor 1 family. Adrenergic receptor subfamily. ADRA2A sub-subfamily.

Protein Information

Name ADRA2A ([HGNC:281](#))

Synonyms ADRA2R, ADRAR

Function Alpha-2 adrenergic receptors are G protein-coupled receptors for catecholamines that activate the G(i/o) protein pathway, thereby promoting adenylyl cyclase inhibition, ERK1/2 stimulation, and voltage-gated calcium channels suppression (PubMed:[2170371](#), PubMed:[23105096](#), PubMed:[2568356](#), PubMed:[35245122](#), PubMed:[27376152](#)). Control a variety of physiological processes, such as regulation of blood pressure, lipolysis and insulin release (PubMed:[2568356](#), PubMed:[27376152](#)). ADRA2A and ADRA2C mediates the presynaptic feedback inhibition of neurotransmitter release from noradrenergic nerve terminals in sympathetic and central nervous systems. ADRA2A inhibits transmitter release at high stimulation frequencies, whereas ADRA2C modulates neurotransmission at lower levels of nerve activity (By similarity). The rank order of potency for agonists of ADRA2A is oxymetazoline > clonidine > epinephrine > norepinephrine > phenylephrine > dopamine > p-synephrine > p-tyramine > serotonin = p-octopamine. For antagonists, the rank order is yohimbine > phentolamine = mianserine > chlorpromazine = spiperone = prazosin > propranolol > alprenolol = pindolol (PubMed:[2170371](#), PubMed:[2568356](#)).


Cellular Location Cell membrane; Multi-pass membrane protein

Background

The alpha-2A adrenergic receptor, also known as ADRA2A denotes the human gene encoding it. This gene is mapped to 10q25.2. Alpha-2-adrenergic receptors are members of the G protein-coupled receptor superfamily. They include 3 highly homologous subtypes: alpha2A, alpha2B, and alpha2C. These receptors have a critical role in regulating neurotransmitter release from sympathetic nerves and from adrenergic neurons in the central nervous system. Studies in mouse revealed that both the alpha2A and alpha2C subtypes were required for normal presynaptic control of transmitter release from sympathetic nerves in the heart and from central noradrenergic neurons; the alpha2A subtype inhibited transmitter release at high stimulation frequencies, whereas the alpha2C subtype modulated neurotransmission at lower levels of nerve activity. This gene encodes alpha2A subtype and it contains no introns in either its coding or untranslated sequences. Alpha-2 adrenergic receptors mediate the catecholamine-induced inhibition of adenylate cyclase through the action of G proteins.

Images

Anti-alpha 2a Adrenergic Receptor antibody, ABO11507,
Western blotting Lane 1: HELA Cell Lysate Lane 2: PANC
Cell Lysate



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.