

Anti-PLTP Antibody

Catalog # ABO11538

Product Information

Application	WB, IHC-P
Primary Accession	P55058
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Phospholipid transfer protein(PLTP) detection. Tested with WB, IHC-P in Human.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	5360
Other Names	Phospholipid transfer protein, Lipid transfer protein II, PLTP
Calculated MW	54739
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, By Heat Western blot, 0.1-0.5 μ g/ml, Human
Subcellular Localization	Secreted.
Tissue Specificity	Wide tissue distribution. Placenta > pancreas > lung > kidney > heart > liver > skeletal muscle > brain.
Source	Eukaryota
Protein Name	Phospholipid transfer protein
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg Na ₃ .
Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human PLTP(256-271aa QLQEEERMVYVAFSEF).
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20° C for one year. After r° Constitution, at 4° C for one month. It° Can also

Sequence Similarities

be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.

Belongs to the BPI/LBP/Plunc superfamily. BPI/LBP family.

Protein Information

Name PLTP

Function Mediates the transfer of phospholipids and free cholesterol from triglyceride-rich lipoproteins (low density lipoproteins or LDL and very low density lipoproteins or VLDL) into high-density lipoproteins (HDL) as well as the exchange of phospholipids between triglyceride-rich lipoproteins themselves (PubMed:[11013307](#), PubMed:[19321130](#), PubMed:[21515415](#), PubMed:[29883800](#), PubMed:[7654777](#), PubMed:[9132017](#)). Facilitates the transfer of a spectrum of different lipid molecules, including diacylglycerol, phosphatidic acid, sphingomyelin, phosphatidylcholine, phosphatidylinositol, phosphatidylglycerol, cerebroside and phosphatidyl ethanolamine (PubMed:[9132017](#)). Plays an important role in HDL remodeling which involves modulating the size and composition of HDL (PubMed:[29883800](#)). Also plays a key role in the uptake of cholesterol from peripheral cells and tissues that is subsequently transported to the liver for degradation and excretion (PubMed:[21736953](#)). Two distinct forms of PLTP exist in plasma: an active form that can transfer phosphatidylcholine from phospholipid vesicles to HDL, and an inactive form that lacks this capability (PubMed:[11013307](#)).


Cellular Location Secreted. Nucleus. Note=Nuclear export is XPO1/CRM1- dependent.

Tissue Location Widely expressed. Highest level of expression in the ovary, thymus and placenta, with moderate levels found in the pancreas, small intestine, testis, lung and prostate. Low level expression in the kidney, liver and spleen, with very low levels found in the heart, colon, skeletal muscle, leukocytes and brain. Expressed in the cortical neurons.


Background

Phospholipid transfer protein (PLTP), also known as lipid transfer protein II is a protein that in humans is encoded by the PLTP gene. This gene is mapped to 20q13.12. The protein encoded by this gene is one of at least two lipid transfer proteins found in human plasma. The encoded protein transfers phospholipids from triglyceride-rich lipoproteins to high density lipoprotein (HDL). In addition to regulating the size of HDL particles, this protein may be involved in cholesterol metabolism. At least two transcript variants encoding different isoforms have been found for this gene.

Images



Anti-PLTP antibody, ABO11538, IHC(P)IHC(P): Human Placenta Tissue



Anti-PLTP antibody, ABO11538, Western blotting All lanes:
 Anti PLTP (ABO11538) at 0.5ug/ml
 Lane 1: MCF-7 Whole Cell Lysate at 40ug
 Lane 2: RAJI Whole Cell Lysate at 40ug
 Lane 3: HELA Whole Cell Lysate at 40ug
 Predicted bind size: 55KD
 Observed bind size: 55KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.