

Anti-Abhd5 Picoband Antibody

Catalog # ABO11639

Product Information

Application	WB, IHC-P
Primary Accession	Q8WTS1
Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for 1-acylglycerol-3-phosphate O-acyltransferase ABHD5(ABHD5) detection. Tested with WB, IHC-P in Human;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	51099
Other Names	1-acylglycerol-3-phosphate O-acyltransferase ABHD5, 2.3.1.51, Abhydrolase domain-containing protein 5, Lipid droplet-binding protein CGI-58, ABHD5, NCIE2
Calculated MW	39096
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat Western blot, 0.1-0.5 µg/ml, Human, Rat
Subcellular Localization	Cytoplasm . Lipid droplet . Colocalized with PLIN and ADRP on the surface of lipid droplets. The localization is dependent upon the metabolic status of the adipocytes and the activity of PKA (By similarity). .
Tissue Specificity	Widely expressed in various tissues, including lymphocytes, liver, skeletal muscle and brain. Expressed by upper epidermal layers and dermal fibroblasts in skin, hepatocytes and neurons (at protein level). .
Source	Eukaryota
Protein Name	1-acylglycerol-3-phosphate O-acyltransferase ABHD5
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	E. coli-derived human Abhd5 recombinant protein (Position: R169-D349). Human Abhd5 shares 96.7% amino acid (aa) sequence identity with both

mouse and rat Abhd5.

Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.

Protein Information


Name	ABHD5 (HGNC:21396)
Synonyms	NCIE2
Function	Coenzyme A-dependent lysophosphatidic acid acyltransferase that catalyzes the transfer of an acyl group on a lysophosphatidic acid (PubMed: 18606822). Functions preferentially with 1-oleoyl- lysophosphatidic acid followed by 1-palmitoyl-lysophosphatidic acid, 1- stearoyl-lysophosphatidic acid and 1-arachidonoyl-lysophosphatidic acid as lipid acceptor. Functions preferentially with arachidonoyl-CoA followed by oleoyl-CoA as acyl group donors (By similarity). Functions in phosphatidic acid biosynthesis (PubMed: 18606822). May regulate the cellular storage of triacylglycerol through activation of the phospholipase PNPLA2 (PubMed: 16679289). Involved in keratinocyte differentiation (PubMed: 18832586). Regulates lipid droplet fusion (By similarity).
Cellular Location	Cytoplasm. Lipid droplet {ECO:0000250 UniProtKB:Q9DBL9}. Cytoplasm, cytosol {ECO:0000250 UniProtKB:Q9DBL9}. Note=Colocalized with PLIN and ADRP on the surface of lipid droplets. The localization is dependent upon the metabolic status of the adipocytes and the activity of PKA (By similarity).
Tissue Location	Widely expressed in various tissues, including lymphocytes, liver, skeletal muscle and brain. Expressed by upper epidermal layers and dermal fibroblasts in skin, hepatocytes and neurons (at protein level).

Background

1-acylglycerol-3-phosphate O-acyltransferase ABHD5 is an enzyme that in humans is encoded by the ABHD5 gene. The protein encoded by this gene belongs to a large family of proteins defined by an alpha/beta hydrolase fold, and contains three sequence motifs that correspond to a catalytic triad found in the esterase/lipase/thioesterase subfamily. It differs from other members of this subfamily in that its putative catalytic triad contains an asparagine instead of the serine residue. Mutations in this gene have been associated with Chanarin-Dorfman syndrome, a triglyceride storage disease with impaired long-chain fatty acid oxidation.

Images

Western blot analysis of Abhd5 expression in rat kidney extract (lane 1) and A431 whole cell lysates (lane 2). Abhd5 at 43KD was detected using rabbit anti- Abhd5 Antigen Affinity purified polyclonal antibody (Catalog # ABO11639) at 0.5 µg/mL. The blot was developed using chemiluminescence (ECL) method .



Abhd5 was detected in paraffin-embedded sections of human intestinal cancer tissues using rabbit anti- Abhd5 Antigen Affinity purified polyclonal antibody (Catalog # ABO11639) at 1 µg/mL. The immunohistochemical section was developed using SABC method .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.