

Anti-ADH4 Picoband Antibody

Catalog # ABO11648

Product Information

Application	WB
Primary Accession	P08319
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Alcohol dehydrogenase 4(ADH4) detection. Tested with WB in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	127
Other Names	Alcohol dehydrogenase 4, 1.1.1.1, Alcohol dehydrogenase class II pi chain, ADH4
Calculated MW	40222
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat
Subcellular Localization	Cytoplasm.
Source	Eukaryota
Protein Name	Alcohol dehydrogenase 4
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	E. coli-derived human ADH4 recombinant protein (Position: K218-F380). Human ADH4 shares 68.7% and 71.8% amino acid (aa) sequence identity with mouse and rat ADH4, respectively.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins.
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.


Protein Information

Name	ADH4 (HGNC:252)
Function	Catalyzes the NAD-dependent oxidation of either all-trans- retinol or 9-cis-retinol (PubMed: 17279314). Also oxidizes long chain omega-hydroxy fatty acids, such as 20-HETE, producing both the intermediate aldehyde, 20-oxoarachidonate and the end product, a dicarboxylic acid, (5Z,8Z,11Z,14Z)-eicosatetraenedioate (PubMed: 16081420). Also catalyzes the reduction of benzoquinones (PubMed: 10514444).
Cellular Location	Cytoplasm.

Background

Alcohol dehydrogenase 4 is an enzyme that in humans is encoded by the ADH4 gene. This gene encodes class II alcohol dehydrogenase 4 pi subunit, which is a member of the alcohol dehydrogenase family. Members of this enzyme family metabolize a wide variety of substrates, including ethanol, retinol, other aliphatic alcohols, hydroxysteroids, and lipid peroxidation products. Class II alcohol dehydrogenase is a homodimer composed of 2 pi subunits. It exhibits a high activity for oxidation of long-chain aliphatic alcohols and aromatic alcohols and is less sensitive to pyrazole. This gene is localized to chromosome 4 in the cluster of alcohol dehydrogenase genes.

Images



Western blot analysis of ADH4 expression in rat liver extract (lane 1), mouse liver extract (lane 2) and HEPG2 whole cell lysates (lane 3). ADH4 at 40KD was detected using rabbit anti- ADH4 Antigen Affinity purified polyclonal antibody (Catalog # ABO11648) at 0.5 µg/mL. The blot was developed using chemiluminescence (ECL) method .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.