

Anti-EPCAM Picoband Antibody

Catalog # ABO11677

Product Information

Application	WB, IHC-P, IHC-F, FC, ICC, E
Primary Accession	P16422
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Epithelial cell adhesion molecule(EPCAM) detection. Tested with WB, IHC-P, IHC-F, ICC, ELISA, FCM in Human.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	4072
Other Names	Epithelial cell adhesion molecule, Ep-CAM, Adenocarcinoma-associated antigen, Cell surface glycoprotein Trop-1, Epithelial cell surface antigen, Epithelial glycoprotein, EGP, Epithelial glycoprotein 314, EGP314, hEGP314, KS 1/4 antigen, KSA, Major gastrointestinal tumor-associated protein GA733-2, Tumor-associated calcium signal transducer 1, CD326, EPCAM, GA733-2, M1S2, M4S1, MIC18, TACSTD1, TROP1
Calculated MW	34932
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, By Heat Immunohistochemistry(Frozen Section), 0.5-1 μ g/ml Immunocytochemistry, 0.5-1 μ g/ml ELISA , 0.1-0.5 μ g/ml Western blot, 0.1-0.5 μ g/ml Flow Cytometry, 1-3 μ g/1x10 ⁶ cells
Subcellular Localization	Lateral cell membrane ; Single-pass type I membrane protein . Cell junction, tight junction . Colocalizes with CLDN7 at the lateral cell membrane and tight junction. .
Tissue Specificity	Highly and selectively expressed by undifferentiated rather than differentiated embryonic stem cells (ESC). Levels rapidly diminish as soon as ESC's differentiate (at protein levels). Expressed in almost all epithelial cell membranes but not on mesodermal or neural cell membranes. Found on the surface of adenocarcinoma. .

Source	Eukaryota
Protein Name	Epithelial cell adhesion molecule
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human EPCAM (147-189aa ELKHKAREKPYDSKSLRTALQKEITTRYQLDPKFITSILYENN), different from the related mouse sequence by fifteen amino acids, and from the related rat sequence by sixteen ami
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.

Protein Information


Name	EPCAM
Synonyms	GA733-2, M1S2, M4S1, MIC18, TACSTD1, TRO
Function	May act as a physical homophilic interaction molecule between intestinal epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium for providing immunological barrier as a first line of defense against mucosal infection. Plays a role in embryonic stem cells proliferation and differentiation. Up-regulates the expression of FABP5, MYC and cyclins A and E.
Cellular Location	Lateral cell membrane; Single-pass type I membrane protein. Cell junction, tight junction. Note=Colocalizes with CLDN7 at the lateral cell membrane and tight junction
Tissue Location	Highly and selectively expressed by undifferentiated rather than differentiated embryonic stem cells (ESC) Levels rapidly diminish as soon as ESC's differentiate (at protein levels). Expressed in almost all epithelial cell membranes but not on mesodermal or neural cell membranes. Found on the surface of adenocarcinoma.

Background


Epithelial cell adhesion molecule (EpcAM) is a transmembrane glycoprotein mediating Ca²⁺-independent homotypic cell-cell adhesion in epithelia. This gene encodes a carcinoma-associated antigen and is a member of a family that includes at least two type I membrane proteins. This antigen is expressed on most normal epithelial cells and gastrointestinal carcinomas and functions as a homotypic calcium-independent cell adhesion molecule. The antigen is being used as a target for immunotherapy treatment of human carcinomas. Mutations in this gene result in congenital tufting enteropathy.

Images


Western blot analysis of EPCAM expression in HELA whole cell lysates (lane 1), A549 whole cell lysates (lane 2) and



PANC-1 whole cell lysates (lane 3). EPCAM at 40KD was detected using rabbit anti- EPCAM Antigen Affinity purified polyclonal antibody (Catalog # ABO11677) at 0.5 μ g/mL. The blot was developed using chemiluminescence (ECL) method .



EPCAM was detected in paraffin-embedded sections of human intestinal cancer tissues using rabbit anti- EPCAM Antigen Affinity purified polyclonal antibody (Catalog # ABO11677) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .



EPCAM was detected in paraffin-embedded sections of human prostatic cancer tissues using rabbit anti- EPCAM Antigen Affinity purified polyclonal antibody (Catalog # ABO11677) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.