

Anti-HPRT Picoband Antibody

Catalog # ABO11689

Product Information

Application	WB
Primary Accession	P00492
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Hypoxanthine-guanine phosphoribosyltransferase(HPRT1) detection. Tested with WB in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	3251
Other Names	Hypoxanthine-guanine phosphoribosyltransferase, HGPRT, HGPRTase, 2.4.2.8, HPRT1, HPRT
Calculated MW	24579
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat
Subcellular Localization	Cytoplasm.
Source	Eukaryota
Protein Name	Hypoxanthine-guanine phosphoribosyltransferase
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E. coli-derived human HPRT recombinant protein (Position: A2-S110). Human HPRT shares 98.1% and 96.3% amino acid (aa) sequence identity with mouse and rat HPRT, respectively.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins.
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.


Protein Information

Name	HPRT1
Synonyms	HPRT
Function	Converts guanine to guanosine monophosphate, and hypoxanthine to inosine monophosphate. Transfers the 5-phosphoribosyl group from 5-phosphoribosylpyrophosphate onto the purine. Plays a central role in the generation of purine nucleotides through the purine salvage pathway.
Cellular Location	Cytoplasm.

Background

Hypoxanthine-guanine phosphoribosyltransferase (HGPRT) is an enzyme encoded in humans by the HPRT1 gene. The protein encoded by this gene is a transferase, which catalyzes conversion of hypoxanthine to inosine monophosphate and guanine to guanosine monophosphate via transfer of the 5-phosphoribosyl group from 5-phosphoribosyl 1-pyrophosphate. This enzyme plays a central role in the generation of purine nucleotides through the purine salvage pathway. Mutations in this gene result in Lesch-Nyhan syndrome or gout.

Images



Western blot analysis of HPRT1 expression in rat brain extract (lane 1), mouse brain extract (lane 2) and HELA whole cell lysates (lane 3). HPRT1 at 24KD was detected using rabbit anti- HPRT1 Antigen Affinity purified polyclonal antibody (Catalog # ABO11689) at 0.5 µg/mL. The blot was developed using chemiluminescence (ECL) method .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.