

Anti-Adiponectin Picoband Antibody

Catalog # ABO11726

Product Information

Application	WB, IHC-P
Primary Accession	Q15848
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Adiponectin(ADIPOQ) detection. Tested with WB, IHC-P in Human.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	9370
Other Names	Adiponectin, 30 kDa adipocyte complement-related protein, Adipocyte complement-related 30 kDa protein, ACRP30, Adipocyte, C1q and collagen domain-containing protein, Adipose most abundant gene transcript 1 protein, apM-1, Gelatin-binding protein, ADIPOQ, ACDC, ACRP30, APM1, GBP28
Calculated MW	26414
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, By Heat Western blot, 0.1-0.5 μ g/ml, Human
Subcellular Localization	Secreted.
Tissue Specificity	Synthesized exclusively by adipocytes and secreted into plasma.
Source	Eukaryota
Protein Name	Adiponectin
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E.coli-derived human adiponectin recombinant protein (Position: E19-N244). Human adiponectin shares 85% amino acid (aa) sequence identity with mouse adiponectin.
Purification	Immunogen affinity purified.

Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Contains 1 C1q domain.


Protein Information

Name	ADIPOQ
Function	Important adipokine involved in the control of fat metabolism and insulin sensitivity, with direct anti-diabetic, anti-atherogenic and anti-inflammatory activities. Stimulates AMPK phosphorylation and activation in the liver and the skeletal muscle, enhancing glucose utilization and fatty-acid combustion. Antagonizes TNF by negatively regulating its expression in various tissues such as liver and macrophages, and also by counteracting its effects. Inhibits endothelial NF-kappa-B signaling through a cAMP-dependent pathway. May play a role in cell growth, angiogenesis and tissue remodeling by binding and sequestering various growth factors with distinct binding affinities, depending on the type of complex, LMW, MMW or HMW.
Cellular Location	Secreted.
Tissue Location	Synthesized exclusively by adipocytes and secreted into plasma.

Background


ADIPOQ (Adipocyte-, C1q-, and Collagen Domain-Containing),also known as APM1, ADPN or ACDC, is a protein which in humans is encoded by the ADIPOQ gene. Using FISH, Das et al. (2001) mapped the mouse Acrp30 gene to chromosome 16 in a region showing homology of synteny with human 3q27. By RNase protection and Western blot analysis, Schaffler et al. (1999) showed that APM1 is expressed by differentiated adipocytes as a 33-kD protein that is also detectable in serum. By sequence comparisons, they found links between APM1 and TNF family ligands as well as to cytokines expressed by T cells. Adiponectin is a protein hormone that modulates a number of metabolic processes, including glucose regulation and fatty acid oxidation. Adiponectin is exclusively secreted from adipose tissue (and also from the placenta in pregnancy) into the bloodstream and is very abundant in plasma relative to many hormones.

Images





Anti- Adiponectin antibody, ABO11726, IHC(P)IHC(P):
Human Lung Cancer Tissue

Anti- Adiponectin antibody, ABO11726, IHC(P)IHC(P):
Human Mammary Cancer Tissue



Anti- Adiponectin antibody, ABO11726, IHC(P)IHC(P):
Human Placenta Tissue



Anti- Adiponectin antibody, ABO11726, Western
blottingAll lanes: Anti Adiponectin (ABO11726) at
0.5ug/mlLane 1: Human Placenta Tissue Lysate at
50ugLane 2: HELA Whole Cell Lysate at 40ugLane 3:
JURKAT Whole Cell Lysate at 40ugPredicted bind size:
30KDObserved bind size: 30KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.