

Anti-BRCA1 Picoband Antibody

Catalog # ABO11740

Product Information

Application	WB, IHC-P
Primary Accession	P38398
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Breast cancer type 1 susceptibility protein(BRCA1) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	672
Other Names	Breast cancer type 1 susceptibility protein, 2.3.2.27, RING finger protein 53, RING-type E3 ubiquitin transferase BRCA1, BRCA1, RNF53
Calculated MW	207721
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5 μ g/ml, Human
Subcellular Localization	Nucleus . Chromosome . Cytoplasm . Localizes at sites of DNA damage at double-strand breaks (DSBs); recruitment to DNA damage sites is mediated by the BRCA1-A complex. Translocated to the cytoplasm during UV-induced apoptosis. .
Tissue Specificity	Isoform 1 and isoform 3 are widely expressed. Isoform 3 is reduced or absent in several breast and ovarian cancer cell lines.
Source	Eukaryota
Protein Name	Breast cancer type 1 susceptibility protein
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E.coli-derived human BRCA1 recombinant protein (Position: E1661-Y1863). Human BRCA1 shares 65% and 66% amino acid (aa) sequences identity with mouse and rat BRCA1, respectively.
Purification	Immunogen affinity purified.

Cross Reactivity	No cross reactivity with other proteins
Storage	At -20° C for one year. After r° Constitution, at 4° C for one month. It° Can also be aliquotted and stored frozen at -20° C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Contains 2 BRCT domains.

Protein Information


Name	BRCA1
Synonyms	RNF53
Function	E3 ubiquitin-protein ligase that specifically mediates the formation of 'Lys-6'-linked polyubiquitin chains and plays a central role in DNA repair by facilitating cellular responses to DNA damage (PubMed: 10500182 , PubMed: 12887909 , PubMed: 12890688 , PubMed: 14976165 , PubMed: 16818604 , PubMed: 17525340 , PubMed: 19261748). It is unclear whether it also mediates the formation of other types of polyubiquitin chains (PubMed: 12890688). The BRCA1-BARD1 heterodimer coordinates a diverse range of cellular pathways such as DNA damage repair, ubiquitination and transcriptional regulation to maintain genomic stability (PubMed: 12890688 , PubMed: 14976165 , PubMed: 20351172). Regulates centrosomal microtubule nucleation (PubMed: 18056443). Required for appropriate cell cycle arrests after ionizing irradiation in both the S-phase and the G2 phase of the cell cycle (PubMed: 10724175 , PubMed: 11836499 , PubMed: 12183412 , PubMed: 19261748). Required for FANCD2 targeting to sites of DNA damage (PubMed: 12887909). Inhibits lipid synthesis by binding to inactive phosphorylated ACACA and preventing its dephosphorylation (PubMed: 16326698). Contributes to homologous recombination repair (HRR) via its direct interaction with PALB2, fine-tunes recombinational repair partly through its modulatory role in the PALB2-dependent loading of BRCA2-RAD51 repair machinery at DNA breaks (PubMed: 19369211). Component of the BRCA1-RBBP8 complex which regulates CHEK1 activation and controls cell cycle G2/M checkpoints on DNA damage via BRCA1-mediated ubiquitination of RBBP8 (PubMed: 16818604). Acts as a transcriptional activator (PubMed: 20160719).
Cellular Location	Nucleus. Chromosome. Cytoplasm. Note=Localizes at sites of DNA damage at double-strand breaks (DSBs); recruitment to DNA damage sites is mediated by ABRAXAS1 and the BRCA1-A complex (PubMed: 26778126) Translocated to the cytoplasm during UV-induced apoptosis (PubMed: 20160719). [Isoform 5]: Cytoplasm
Tissue Location	Isoform 1 and isoform 3 are widely expressed. Isoform 3 is reduced or absent in several breast and ovarian cancer cell lines

Background


BRCA1, also known as BRCC1, is a gene which mapping to 17q21.3. This gene encodes a nuclear phosphoprotein that plays a role in maintaining genomic stability, and it also acts as a tumor suppressor. The encoded protein combines with other tumor suppressors, DNA damage sensors, and signal transducers to form a large multi-subunit protein complex known as the BRCA1-associated genome surveillance complex (BASC). BRCA1 product associates with RNA polymerase II, and through the C-terminal domain, also interacts with histone deacetylase complexes. This protein thus plays a role in transcription, DNA repair of double-stranded breaks, and recombination. In addition to it, BRCA1 may normally serve as a negative regulator of mammary epithelial cell growth and that this function is compromised in breast cancer either

by direct mutation or by alterations in gene expression.


Images




Anti-BRCA1 Picoband antibody, ABO11740-1.JPGIHC(P):
Mouse Testis Tissue



Anti-BRCA1 Picoband antibody, ABO11740-2.JPGIHC(P):
Rat Testis Tissue



Anti-BRCA1 Picoband antibody, ABO11740-3.JPGIHC(P):
Human Mammary Cancer Tissue



Anti-BRCA1 Picoband antibody, ABO11740-4.jpgAll lanes:
Anti-BRCA1(ABO11740) at 0.5ug/mlLane 1: HELA Whole
Cell Lysate at 40ugLane 2: MCF-7 Whole Cell Lysate at
40ugLane 3: A549 Whole Cell Lysate at 40ugPredicted
bind size: 207KDObserved bind size: 207KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.