

Anti-ICAM-1 Antibody

Catalog # ABO11742

Product Information

Application	WB, IHC-P
Primary Accession	P05362
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Intercellular adhesion molecule 1(ICAM1) detection. Tested with WB, IHC-P in Human.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	3383
Other Names	Intercellular adhesion molecule 1, ICAM-1, Major group rhinovirus receptor, CD54, ICAM1
Calculated MW	57825
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat Western blot, 0.1-0.5 µg/ml, Human
Subcellular Localization	Membrane; Single-pass type I membrane protein.
Source	Eukaryota
Protein Name	Intercellular adhesion molecule 1
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E.coli-derived human ICAM1 recombinant protein (Position: L214-P532). Human ICAM1 shares 52% and 48% amino acid (aa) sequences identity with mouse and rat ICAM1, respectively.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.


Protein Information

Name	ICAM1 (HGNC:5344)
Function	Cell adhesion molecule that functions as a receptor ligand of the signaling receptor ITGAL:ITGB2/LFA-1 (lymphocyte-function associated (LFA) molecule 1) ensuring leukocyte cell-cell adhesion, by providing a calibrated system to namely adjust T-cell killing to the antigen stimulation strength (PubMed: 3086451 , PubMed: 3340213 , PubMed: 38195629). Also functions as a ligand receptor of the signaling receptor ITGAM:ITGB2/MAC-1 ensuring adhesion between stimulated neutrophils and stimulated endothelial cells (PubMed: 1980124). During leukocyte trans-endothelial migration, ICAM1 engagement promotes the assembly of endothelial apical cups through ARHGEF26/SGEF and RHOG activation (PubMed: 17875742). Promotes cell aggregation in epithelial cells through interaction with MUC1 (PubMed: 11173916).
Cellular Location	Cell membrane; Single-pass type I membrane protein
Tissue Location	Expressed on non-hematopoietic cells such as vascular endothelial cells, thymic epithelial cells, certain other epithelial cells, and fibroblasts, and on hematopoietic cells such as tissue macrophages, mitogen-stimulated T lymphocyte blasts, and germinal center dendritic cells in tonsils, lymph nodes, and Peyer's patches (PubMed:3086451). Expressed in low amounts on peripheral blood leukocytes (PubMed:3086451).

Background


Intercellular adhesion molecule-1 (ICAM-1) is an integral membrane protein, a member of the immunoglobulin superfamily, and a ligand for lymphocyte function-associated (LFA) antigens, a beta 2 leukocyte integrin. The normal function of human ICAM-1 is to provide adhesion between endothelial cells and leukocytes after injury or stress. ICAM-1 binds to leukocyte function-associated antigen (LFA-1) or macrophage-1 antigen (Mac-1). It is found on leukocytes, fibroblasts, epithelial cells and endothelial cells and its expression is regulated by inflammatory cytokines. ICAM-1 has a tissue distribution similar to that of the major histocompatibility complex class II antigens and is likely to play a role in inflammatory responses.

Images



Anti-ICAM1 Picoband antibody, ABO11742-1.JPGIHC(P):
Human Intestinal Cancer Tissue

Anti-ICAM1 Picoband antibody, ABO11742-2.jpgAll lanes:
Anti-ICAM1(ABO11742) at 0.5ug/mlLane 1: HEPG2 Whole
Cell Lysate at 40ugLane 2: K562 Whole Cell Lysate at
40ugLane 3: A549 Whole Cell Lysate at 40ugLane 4:
HT1080 Whole Cell Lysate at 40ugPredicted bind size:



58KDObserved bind size: 100KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.