

Anti-CCL17/TARC Antibody

Catalog # ABO11755

Product Information

Application	WB, E
Primary Accession	Q9WUJ6
Host	Rabbit
Reactivity	Mouse
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for C-C motif chemokine 17(CCL17) detection. Tested with WB, ELISA in Mouse.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	20295
Calculated MW	10467
Application Details	Western blot, 0.1-0.5 µg/ml, Mouse, - ELISA , 0.1-0.5 µg/ml, Mouse
Subcellular Localization	Secreted .
Source	Eukaryota
Protein Name	C-C motif chemokine 17
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	E.coli-derived mouse TARC recombinant protein (Position: A34-P103). Mouse TARC shares 71% amino acid (aa) sequence identity with human TARC.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.

Protein Information

Name	Ccl17
-------------	-------

Synonyms	Tarc {ECO:0000303 PubMed:10508243}
Function	Chemokine, which displays chemotactic activity for T lymphocytes, preferentially Th2 cells, but not monocytes or granulocytes. Therefore plays an important role in a wide range of inflammatory and immunological processes (PubMed: 10508243 , PubMed: 10508268). Acts by binding to CCR4 at T-cell surface (By similarity). Mediates GM-CSF/CSF2-driven pain and inflammation (PubMed: 27525438). In the brain, required to maintain the typical, highly branched morphology of hippocampal microglia under homeostatic conditions. May be important for the appropriate adaptation of microglial morphology and synaptic plasticity to acute lipopolysaccharide (LPS)-induced neuroinflammation (PubMed: 30277599). Plays a role in wound healing, mainly by inducing fibroblast migration into the wound (PubMed: 21521373).
Cellular Location	Secreted
Tissue Location	Expressed in primary and secondary lymphoid organs, where it is specifically produced by a subset of dendritic cells. Not expressed in the spleen (PubMed:12615900). Constitutively expressed in thymus, as well as in the lung, skin and intestine (PubMed:10508243, PubMed:10508268, PubMed:12615900). Not expressed in bone marrow-derived macrophages and activated B cells, nor in thymocytes (PubMed:10508243, PubMed:10508268). In the brain, predominantly expressed in a subset of hippocampal CA1 neurons (PubMed:30277599)

Background

Chemokine (C-C motif) ligand 17 (CCL17) is a small cytokine belonging to the CC chemokine family that is also known as thymus and activation regulated chemokine (TARC). CCL17 is expressed constitutively in thymus, but only transiently in phytohemagglutinin-stimulated peripheral blood mononuclear cells. This chemokine specifically binds and induces chemotaxis in T cells and elicits its effects by interacting with the chemokine receptor CCR4. The gene for CCL17 is located on chromosome 16, in humans, along with other chemokines called CCL22 and CX3CL1. The standard used in this kit is recombinant human CCL17, consisting of 71 amino acids with the molecular weight of 8Kda.

Images



Anti-TARC Picoband antibody, ABO11755-1.jpg
 All lanes:
 Anti-TARC(ABO11755) at 0.5ug/ml
 WB: Recombinant Mouse TARC Protein 0.5ng
 Predicted bind size: 34KD
 Observed bind size: 34KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.