

Anti-Wnt5a Picoband Antibody

Catalog # ABO11779

Product Information

Application	WB, IHC-P
Primary Accession	P41221
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Protein Wnt-5a(WNT5A) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	7474
Other Names	Protein Wnt-5a, WNT5A
Calculated MW	42339
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse
Subcellular Localization	Secreted, extracellular space, extracellular matrix.
Tissue Specificity	Expression is increased in differentiated thyroid carcinomas compared to normal thyroid tissue and anaplastic thyroid tumors where expression is low or undetectable. Expression is found in thyrocytes but not in stromal cells (at protein level). .
Source	Eukaryota
Protein Name	Protein Wnt-5a
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E.coli-derived human Wnt5a recombinant protein (Position: D184-D303). Human Wnt5a shares 100% amino acid (aa) sequence identity with both mouse and rat Wnt5a.
Purification	Immunogen affinity purified.

Cross Reactivity	No cross reactivity with other proteins
Storage	At -20° C for one year. After r° Constitution, at 4° C for one month. It° Can also be aliquotted and stored frozen at -20° C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the Wnt family.

Protein Information



Name	WNT5A
Function	Ligand for members of the frizzled family of seven transmembrane receptors. Can activate or inhibit canonical Wnt signaling, depending on receptor context. In the presence of FZD4, activates beta-catenin signaling. In the presence of ROR2, inhibits the canonical Wnt pathway by promoting beta-catenin degradation through a GSK3-independent pathway which involves down-regulation of beta- catenin-induced reporter gene expression (By similarity). Suppression of the canonical pathway allows chondrogenesis to occur and inhibits tumor formation. Stimulates cell migration. Decreases proliferation, migration, invasiveness and clonogenicity of carcinoma cells and may act as a tumor suppressor (PubMed: 15735754). Mediates motility of melanoma cells (PubMed: 17426020). Required during embryogenesis for extension of the primary anterior-posterior axis and for outgrowth of limbs and the genital tubercle. Inhibits type II collagen expression in chondrocytes (By similarity).
Cellular Location	Secreted, extracellular space, extracellular matrix. Secreted
Tissue Location	Expression is increased in differentiated thyroid carcinomas compared to normal thyroid tissue and anaplastic thyroid tumors where expression is low or undetectable. Expression is found in thyrocytes but not in stromal cells (at protein level) (PubMed:15735754). Detected in neonate heart and lung (PubMed:8288227)

Background

Protein Wnt-5a is a protein that in humans is encoded by the WNT5A gene. This gene is a member of the WNT gene family. The WNTs comprise a large class of secreted proteins that control essential developmental processes such as embryonic patterning, cell growth, migration, and differentiation. It is mapped to 3p14.3. WNT5A is highly expressed in the dermal papilla of depilated skin. This protein plays an essential role in regulating developmental pathways during embryogenesis. And this protein may also play a role in oncogenesis. WNT5A and DVL can promote axon differentiation mediated by the PAR3-PAR6-aPKC complex.

Images

Anti-Wnt5a Picoband antibody, ABO11779-1.jpgAll lanes:
Anti-WNT5A(ABO11779) at 0.5ug/mlLane 1: HELA Whole
Cell Lysate at 40ugLane 2: MCF-7 Whole Cell Lysate at
40ugLane 3: HT1080 Whole Cell Lysate at 40ugPredicted
bind size: 41KDObserved bind size: 41KD



IHC(P): Human Placenta Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.