

Anti-CBL Picoband Antibody

Catalog # ABO11791

Product Information

Application	WB
Primary Accession	P22681
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for E3 ubiquitin-protein ligase CBL(CBL) detection. Tested with WB in Human.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	867
Other Names	E3 ubiquitin-protein ligase CBL, 2.3.2.27, Casitas B-lineage lymphoma proto-oncogene, Proto-oncogene c-Cbl, RING finger protein 55, RING-type E3 ubiquitin transferase CBL, Signal transduction protein CBL, CBL, CBL2, RNF55
Calculated MW	99633
Application Details	Western blot, 0.1-0.5 µg/ml, Human
Subcellular Localization	Cytoplasm. Cell membrane. Colocalizes with FGFR2 in lipid rafts at the cell membrane.
Source	Eukaryota
Protein Name	E3 ubiquitin-protein ligase CBL
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E.coli-derived human CBL recombinant protein (Position: A556-T906). Human CBL shares 84% amino acid (aa) sequence identity with mouse CBL.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.
Sequence Similarities	Contains 1 Cbl-PTB (Cbl-type phosphotyrosine-binding) domain.


Protein Information

Name	CBL
Synonyms	CBL2, RNF55
Function	E3 ubiquitin-protein ligase that acts as a negative regulator of many signaling pathways by mediating ubiquitination of cell surface receptors (PubMed:10514377, PubMed:11896602, PubMed:14661060, PubMed:14739300, PubMed:15190072, PubMed:17509076, PubMed:18374639, PubMed:19689429, PubMed:21596750, PubMed:28381567, PubMed:40101708). Accepts ubiquitin from specific E2 ubiquitin-conjugating enzymes, and then transfers it to substrates promoting their degradation by the proteasome (PubMed:10514377, PubMed:14661060, PubMed:14739300, PubMed:17094949, PubMed:17509076, PubMed:17974561). Recognizes activated receptor tyrosine kinases, including KIT, FLT1, FGFR1, FGFR2, PDGFRA, PDGFRB, CSF1R, EPHA8 and KDR and mediates their ubiquitination to terminate signaling (PubMed:15190072, PubMed:18374639, PubMed:21596750). Recognizes membrane-bound HCK, SRC and other kinases of the SRC family and mediates their ubiquitination and degradation (PubMed:11896602). Ubiquitinates EGFR and SPRY2 (PubMed:17094949, PubMed:17974561). Involved in LAG3-mediated inhibition of TCR signaling: following ligand-binding to LAG3, catalyzes 'Lys-63'-linked ubiquitination of LAG3, unleashing the LAG3 C-terminus from the membrane, and initiating a signaling that prevents TCR activation (PubMed:40101708). Ubiquitinates NECTIN1 following association between NECTIN1 and herpes simplex virus 1/HHV-1 envelope glycoprotein D, leading to NECTIN1 removal from cell surface (PubMed:28381567). Participates in signal transduction in hematopoietic cells. Plays an important role in the regulation of osteoblast differentiation and apoptosis (PubMed:15190072, PubMed:18374639). Essential for osteoclastic bone resorption (PubMed:14739300). The 'Tyr-731' phosphorylated form induces the activation and recruitment of phosphatidylinositol 3-kinase to the cell membrane in a signaling pathway that is critical for osteoclast function (PubMed:14739300). In association with CBLB, required for proper feedback inhibition of ciliary platelet-derived growth factor receptor-alpha (PDGFRA) signaling pathway via ubiquitination and internalization of PDGFRA (By similarity).
Cellular Location	Cytoplasm. Cell membrane. Cell projection, cilium. Golgi apparatus. Note=Colocalizes with FGFR2 in lipid rafts at the cell membrane

Background

CBL(Cbl proto-oncogene) is also known as C-CBL, RNF55, CBL2 and E3 ubiquitin protein ligase. CBL is mapped to chromosome 11q23.3-qter by molecular characterization of the breakpoints in 2 somatic cell hybrids. The encoded protein is one of the enzymes required for targeting substrates for degradation by the proteasome. This protein mediates the transfer of ubiquitin from ubiquitin conjugating enzymes(E2) to specific substrates. This protein also contains an N-terminal phosphotyrosine binding domain that allows it to interact with numerous tyrosine-phosphorylated substrates and target them for proteasome degradation. As such it functions as a negative regulator of many signal transduction pathways. This gene has been found to be mutated or translocated in many cancers including acute myeloid leukaemia. Mutations in this gene are also the cause of Noonan syndrome-like disorder.

Images



Anti-CBL Picoband antibody, ABO11791-1.jpg
 All lanes:
 Anti-CBL(ABO11791) at 0.5ug/ml
 Lane 1: Hela Whole Cell Lysate at 40ug
 Lane 2: MCF-7 Whole Cell Lysate at 40ug
 Lane 3: HepG2 Whole Cell Lysate at 40ug
 Lane 4: Colo320 Whole Cell Lysate at 40ug
 Lane 5: PANC Whole Cell Lysate at 40ug
 Lane 6: SW620 Whole Cell Lysate at 40ug
 Lane 7: A549 Whole Cell Lysate at 40ug
 Lane 8: Skov Whole Cell Lysate at 40ug
 Lane 9: HT1080 Whole Cell Lysate at 40ug
 Predicted bind size: 100KD
 Observed bind size: 100KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.