

Anti-RIP Picoband Antibody

Catalog # ABO11820

Product Information

Application	WB
Primary Accession	Q13546
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Receptor-interacting serine/threonine-protein kinase 1(RIPK1) detection. Tested with WB in Human.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	8737
Other Names	Receptor-interacting serine/threonine-protein kinase 1, 2.7.11.1, Cell death protein RIP, Receptor-interacting protein 1, RIP-1, Serine/threonine-protein kinase RIP, RIPK1, RIP, RIP1
Calculated MW	75931
Application Details	Western blot, 0.1-0.5 µg/ml, Human
Subcellular Localization	Cytoplasm. Cell membrane .
Source	Eukaryota
Protein Name	Receptor-interacting serine/threonine-protein kinase 1
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	E.coli-derived human RIP recombinant protein (Position: K316-N671). Human RIP shares 65% amino acid (aa) sequence identity with mouse RIP.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family.


Protein Information

Name	RIPK1 (HGNC:10019)
Function	Serine-threonine kinase which is a key regulator of TNF- mediated apoptosis, necroptosis and inflammatory pathways (PubMed:17703191, PubMed:24144979, PubMed:31827280, PubMed:31827281, PubMed:32657447, PubMed:35831301). Exhibits kinase activity-dependent functions that regulate cell death and kinase-independent scaffold functions regulating inflammatory signaling and cell survival (PubMed:11101870, PubMed:19524512, PubMed:19524513, PubMed:29440439, PubMed:30988283). Has kinase-independent scaffold functions: upon binding of TNF to TNFR1, RIPK1 is recruited to the TNF-R1 signaling complex (TNF-RSC also known as complex I) where it acts as a scaffold protein promoting cell survival, in part, by activating the canonical NF-kappa-B pathway (By similarity). Kinase activity is essential to regulate necroptosis and apoptosis, two parallel forms of cell death: upon activation of its protein kinase activity, regulates assembly of two death-inducing complexes, namely complex IIa (RIPK1-FADD-CASP8), which drives apoptosis, and the complex IIb (RIPK1-RIPK3-MLKL), which drives necroptosis (By similarity). RIPK1 is required to limit CASP8- dependent TNFR1-induced apoptosis (By similarity). In normal conditions, RIPK1 acts as an inhibitor of RIPK3-dependent necroptosis, a process mediated by RIPK3 component of complex IIb, which catalyzes phosphorylation of MLKL upon induction by ZBP1 (PubMed:19524512, PubMed:19524513, PubMed:29440439, PubMed:30988283). Inhibits RIPK3- mediated necroptosis via FADD-mediated recruitment of CASP8, which cleaves RIPK1 and limits TNF-induced necroptosis (PubMed:19524512, PubMed:19524513, PubMed:29440439, PubMed:30988283). Required to inhibit apoptosis and necroptosis during embryonic development: acts by preventing the interaction of TRADD with FADD thereby limiting aberrant activation of CASP8 (By similarity). In addition to apoptosis and necroptosis, also involved in inflammatory response by promoting transcriptional production of pro-inflammatory cytokines, such as interleukin-6 (IL6) (PubMed:31827280, PubMed:31827281). Phosphorylates RIPK3: RIPK1 and RIPK3 undergo reciprocal auto- and trans- phosphorylation (PubMed:19524513). Phosphorylates DAB2IP at 'Ser-728' in a TNF-dependent manner, and thereby activates the MAP3K5-JNK apoptotic cascade (PubMed:15310755, PubMed:17389591). Required for ZBP1-induced NF-kappa-B activation in response to DNA damage (By similarity).
Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:Q60855}. Cell membrane {ECO:0000250 UniProtKB:Q9ZUF4}


Background

RIPK1, also known as RIP or RIP1, is an enzyme that in humans is encoded by the RIPK1 gene. It is mapped to 6p25.2. RIPK1 is a key signaling molecule in the programmed necrosis pathway, which plays important roles in development, tissue damage response, and antiviral immunity. RIPK1 is known to have function in a variety of cellular pathways including the NF- κ B pathway and programmed necrotic cell death (necroptosis). The kinase domain, while important for necroptotic (programmed necrotic) functions, it appears dispensable for other lethal, as well as pro-survival roles. Also, proteolytic processing of RIPK1, through both caspase-dependent and -independent mechanisms, triggers lethality that is dependent on the generation of one or more specific C-terminal cleavage product(s) of RIPK1 upon stress.

Images



Anti-RIP Picoband antibody, ABO11820-1.jpg All lanes:
 Anti RIP (ABO11820) at 0.5ug/ml WB: Recombinant
 Human RIP Protein 0.5ng Predicted bind size:
 38KD Observed bind size: 38KD



Anti-RIP Picoband antibody, ABO11820-2.jpg All lanes:
 Anti RIP (ABO11820) at 0.5ug/ml Lane 1: JURKAT Whole
 Cell Lysate at 40ug Lane 2: 22RV1 Whole Cell Lysate at
 40ug Lane 3: MCF-7 Whole Cell Lysate at 40ug Lane 4:
 HELA Whole Cell Lysate at 40ug Lane 5: A549 Whole Cell
 Lysate at 40ug Predicted bind size: 76KD Observed bind
 size: 76KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.