

Anti-Angiopoietin-1 Antibody

Catalog # ABO11827

Product Information

Application	WB
Primary Accession	Q15389
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Angiopoietin-1(ANGPT1) detection. Tested with WB in Human.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	284
Other Names	Angiopoietin-1, ANG-1, ANGPT1, KIAA0003
Calculated MW	57513
Application Details	Western blot, 0.1-0.5 µg/ml, Human
Subcellular Localization	Secreted.
Source	Eukaryota
Protein Name	Angiopoietin-1
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	E.coli-derived human Angiopoietin 1 recombinant protein (Position: H16-N350). Human Angiopoietin 1 shares 96% amino acid (aa) sequence with both mouse and rat Angiopoietin 1.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.
Sequence Similarities	Contains 1 fibrinogen C-terminal domain.


Protein Information

Name	ANGPT1
Synonyms	KIAA0003
Function	Binds and activates TEK/TIE2 receptor by inducing its dimerization and tyrosine phosphorylation. Plays an important role in the regulation of angiogenesis, endothelial cell survival, proliferation, migration, adhesion and cell spreading, reorganization of the actin cytoskeleton, but also maintenance of vascular quiescence. Required for normal angiogenesis and heart development during embryogenesis. After birth, activates or inhibits angiogenesis, depending on the context. Inhibits angiogenesis and promotes vascular stability in quiescent vessels, where endothelial cells have tight contacts. In quiescent vessels, ANGPT1 oligomers recruit TEK to cell-cell contacts, forming complexes with TEK molecules from adjoining cells, and this leads to preferential activation of phosphatidylinositol 3-kinase and the AKT1 signaling cascades. In migrating endothelial cells that lack cell-cell adhesions, ANGPT1 recruits TEK to contacts with the extracellular matrix, leading to the formation of focal adhesion complexes, activation of PTK2/FAK and of the downstream kinases MAPK1/ERK2 and MAPK3/ERK1, and ultimately to the stimulation of sprouting angiogenesis. Mediates blood vessel maturation/stability. Implicated in endothelial developmental processes later and distinct from that of VEGF. Appears to play a crucial role in mediating reciprocal interactions between the endothelium and surrounding matrix and mesenchyme.
Cellular Location	Secreted.


Background

Angiopoietin 1 is a type of angiopoietin and is encoded by the gene ANGPT1. Angiopoietins are proteins with important roles in vascular development and angiogenesis. All angiopoietins bind with similar affinity to an endothelial cell-specific tyrosine-protein kinase receptor. Angiopoietin 1 is mapped to 8q23.1. The protein encoded by this gene is a secreted glycoprotein that activates the receptor by inducing its tyrosine phosphorylation. It plays a critical role in mediating reciprocal interactions between the endothelium and surrounding matrix and mesenchyme. The protein also contributes to blood vessel maturation and stability, and may be involved in early development of the heart. Angiopoietin-1 seems to play a crucial role in mediating reciprocal interactions between the endothelium and surrounding matrix and mesenchyme.

Images



Anti-Angiopoietin 1 Picoband antibody,
ABO11827-1.jpg All lanes: Anti Angiopoietin1 (ABO11827)
at 0.5ug/ml WB: Recombinant Human Angiopoietin1
Protein 0.5ng Predicted bind size: 39KD Observed bind
size: 39KD



Anti-Angiopoietin 1 Picoband antibody,
 ABO11827-2.jpg All lanes: Anti Angiopoietin1 (ABO11827)
 at 0.5ug/ml Lane 1: HELA Whole Cell Lysate at 40ug Lane 2:
 MCF-7 Whole Cell Lysate at 40ug Lane 3: COLO320 Whole
 Cell Lysate at 40ug Lane 4: A549 Whole Cell Lysate at
 40ug Lane 5: HEPG2 Whole Cell Lysate at 40ug Lane 6:
 293T Whole Cell Lysate at 40ug Lane 7: SW620 Whole Cell
 Lysate at 40ug Predicted bind size: 57KD Observed bind
 size: 65KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.