

Anti-SynCAM Picoband Antibody

Catalog # ABO11837

Product Information

Application	WB
Primary Accession	Q9BY67
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Cell adhesion molecule 1(CADM1) detection. Tested with WB in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	23705
Other Names	Cell adhesion molecule 1, Immunoglobulin superfamily member 4, IgSF4, Nectin-like protein 2, NECL-2, Spermatogenic immunoglobulin superfamily, SgIgSF, Synaptic cell adhesion molecule, SynCAM, Tumor suppressor in lung cancer 1, TSLC-1, CADM1 (HGNC:5951)
Calculated MW	48509
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat
Subcellular Localization	Cell membrane ; Single-pass type I membrane protein . Cell junction, synapse . Associates with perinuclear and plasma membranes in vivo. Localized to the basolateral plasma membrane of epithelial cells in gall bladder. .
Source	Eukaryota
Protein Name	Cell adhesion molecule 1
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	E.coli-derived human SynCAM recombinant protein (Position: Q45-P340). Human SynCAM shares 99% amino acid (aa) sequence identity with mouse SynCAM.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins

Storage	At -20 °C for one year. After r° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the nectin family.

Protein Information

Name	CADM1 (HGNC:5951)
Function	Mediates homophilic cell-cell adhesion in a Ca(2+)- independent manner (PubMed: 12050160 , PubMed: 22438059). Also mediates heterophilic cell-cell adhesion with CADM3 and NECTIN3 in a Ca(2+)- independent manner (By similarity). Interaction with CRTAM promotes natural killer (NK) cell cytotoxicity and interferon-gamma (IFN-gamma) secretion by CD8+ cells in vitro as well as NK cell-mediated rejection of tumors expressing CADM1 in vivo (PubMed: 15811952). In mast cells, may mediate attachment to and promote communication with nerves (PubMed: 15905536). CADM1, together with MITF, is essential for development and survival of mast cells in vivo (PubMed: 22438059). By interacting with CRTAM and thus promoting the adhesion between CD8+ T- cells and CD8+ dendritic cells, regulates the retention of activated CD8+ T-cell within the draining lymph node (By similarity). Required for the intestinal retention of intraepithelial CD4+ CD8+ T-cells and, to a lesser extent, intraepithelial and lamina propria CD8+ T-cells and CD4+ T-cells (By similarity). Interaction with CRTAM promotes the adhesion to gut-associated CD103+ dendritic cells, which may facilitate the expression of gut-homing and adhesion molecules on T-cells and the conversion of CD4+ T-cells into CD4+ CD8+ T-cells (By similarity). Acts as a synaptic cell adhesion molecule and plays a role in the formation of dendritic spines and in synapse assembly (By similarity). May be involved in neuronal migration, axon growth, pathfinding, and fasciculation on the axons of differentiating neurons (By similarity). May play diverse roles in the spermatogenesis including in the adhesion of spermatocytes and spermatids to Sertoli cells and for their normal differentiation into mature spermatozoa (By similarity). Acts as a tumor suppressor in non-small-cell lung cancer (NSCLC) cells (PubMed: 11279526 , PubMed: 12234973). May contribute to the less invasive phenotypes of lepidic growth tumor cells (PubMed: 12920246).
Cellular Location	Cell membrane; Single-pass type I membrane protein. Synapse {ECO:0000250 UniProtKB:Q8R5M8} Note=Localized to the basolateral plasma membrane of epithelial cells in gall bladder. {ECO:0000250 UniProtKB:Q8R5M8}

Background

Cell adhesion molecule 1 is a protein that, in humans, is encoded by the CADM1 gene. This gene is also known as IGSF4, SYNCAM or TSLC1, and it is mapped to 11q23.3. CADM1 functions as a homophilic cell adhesion molecule at the synapse. Expression of the isolated cytoplasmic tail of CADM1 in neurons inhibited synapse assembly. Conversely, expression of full-length CADM1 in nonneuronal cells induced synapse formation by cocultured hippocampal neurons with normal release properties.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.