

Anti-CYP2E1 Picoband Antibody

Catalog # ABO11881

Product Information

Application	WB, IHC-P
Primary Accession	P05181
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Cytochrome P450 2E1(CYP2E1) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	1571
Other Names	Cytochrome P450 2E1, 1.14.13.-, 4-nitrophenol 2-hydroxylase, 1.14.13.n7, CYP11E1, Cytochrome P450-J, Cytochrome P450 2E1, N-terminally processed, CYP2E1, CYP2E
Calculated MW	56849
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Mouse, Rat, Human, By Heat Western blot, 0.1-0.5 µg/ml, Human, Rat
Subcellular Localization	Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein.
Source	Eukaryota
Protein Name	Cytochrome P450 2E1
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E.coli-derived human CYP2E1 recombinant protein (Position: M1-Y310). Human CYP2E1 shares 73% and 74% amino acid (aa) sequences identity with mouse and rat CYP2E1, respectively.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20° C for one year. After reconstitution, at 4° C for one month. It can also

Sequence Similarities

be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
Belongs to the cytochrome P450 family.

Protein Information

Name CYP2E1 {ECO:0000303 | PubMed:10553002, ECO:0000312 | HGNC:HGNC:2631}


Function A cytochrome P450 monooxygenase involved in the metabolism of fatty acids (PubMed:[10553002](#), PubMed:[18577768](#)). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed:[10553002](#), PubMed:[18577768](#)). Catalyzes the hydroxylation of carbon-hydrogen bonds. Hydroxylates fatty acids specifically at the omega-1 position displaying the highest catalytic activity for saturated fatty acids (PubMed:[10553002](#), PubMed:[18577768](#)). May be involved in the oxidative metabolism of xenobiotics (Probable).

Cellular Location Endoplasmic reticulum membrane {ECO:0000250 | UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250 | UniProtKB:P05182}. Microsome membrane {ECO:0000250 | UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250 | UniProtKB:P05182}. Mitochondrion inner membrane {ECO:0000250 | UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250 | UniProtKB:P05182}. Note=Post-translationally targeted to mitochondria. TOMM70 is required for the translocation across the mitochondrial outer membrane. After translocation into the matrix, associates with the inner membrane as a membrane extrinsic protein {ECO:0000250 | UniProtKB:P05182}


Background

Cytochrome P450 2E1 (abbreviated CYP2E1), a member of the cytochrome P450 mixed-function oxidase system, is involved in the metabolism of xenobiotics in the body. In humans, the CYP2E1 enzyme is encoded by the CYP2E1 gene. It is mapped to 10q26.3. While it is involved in the oxidative metabolism of a small range of substrates (mostly small polar molecules), there are many important drug interactions mediated by CYP2E1. Most drugs undergo deactivation by CYP2E1, either directly or by facilitated excretion from the body. Also, many substances are bioactivated by CYP2E1 to form their active compounds. In addition, CYP2E1 is an important enzyme for the conversion of ethanol to acetaldehyde and to acetate in humans. In the conversion sequence of acetyl-CoA to glucose, CYP2E1 transforms acetone via acetol into propylene glycol and methylglyoxal, the precursors of pyruvate, acetate and lactate.

Images



Anti- CYP2E1 antibody, ABO11881, IHC(P)IHC(P): Mouse Liver Tissue



Anti- CYP2E1 antibody, ABO11881, IHC(P)IHC(P): Rat Liver Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.