10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-FOXM1 Picoband Antibody

Catalog # ABO11885

Product Information

Application	WB
Primary Accession	Q08050
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Forkhead box protein M1(FOXM1) detection. Tested with WB in Human.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	2305
Other Names	Forkhead box protein M1, Forkhead-related protein FKHL16, Hepatocyte nuclear factor 3 forkhead homolog 11, HFH-11, HNF-3/fork-head homolog 11, M-phase phosphoprotein 2, MPM-2 reactive phosphoprotein 2, Transcription factor Trident, Winged-helix factor from INS-1 cells, FOXM1, FKHL16, HFH11, MPP2, WIN
Calculated MW	84283
Application Details	Western blot, 0.1-0.5 □g/ml, Human
Subcellular Localization	Nucleus.
Tissue Specificity	Expressed in thymus, testis, small intestine, colon followed by ovary. Appears to be expressed only in adult organs containing proliferating/cycling cells or in response to growth factors. Also expressed in epithelial cell lines derived from tumors. Not expressed in resting cells. Isoform 2 is highly expressed in testis.
Source	Eukaryota
Protein Name	Forkhead box protein M1
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	E.coli-derived human FOXM1 recombinant protein (Position: A411-Q763). Human FOXM1 shares 75% and 76% amino acid (aa) sequences identity with mouse and rat FOXM1, respectively.

Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20˚C for one year. After r˚Constitution, at 4˚C for one month. It˚Can also be aliquotted and stored frozen at -20˚C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Contains 1 fork-head DNA-binding domain.


Protein Information

Name	FOXM1
Synonyms	FKHL16, HFH11, MPP2, WIN
Function	Transcription factor regulating the expression of cell cycle genes essential for DNA replication and mitosis (PubMed:19160488, PubMed:20360045). Plays a role in the control of cell proliferation (PubMed:19160488). Also plays a role in DNA break repair, participating in the DNA damage checkpoint response (PubMed:17101782). Promotes transcription of PHB2 (PubMed:33754036).
Cellular Location	[Isoform 1]: Nucleus [Isoform 4]: Cytoplasm
Tissue Location	Expressed in thymus, testis, small intestine, colon followed by ovary. Appears to be expressed only in adult organs containing proliferating/cycling cells or in response to growth factors. Also expressed in epithelial cell lines derived from tumors Not expressed in resting cells. Isoform 2 is highly expressed in testis

Background

Forkhead box protein M1, also called FOXM1, is a protein that in humans is encoded by the FOXM1 gene. It is mapped to 12p13.33. The protein encoded by this gene is a member of the FOX family of transcription factors. FOXM1 is known to play a key role in cell cycle progression where endogenous FOXM1 expression peaks at S and G2/M phases and also in the control of cell proliferation. FOXM1 gene is now known as a human proto-oncogene. Abnormal upregulation of FOXM1 is involved in the oncogenesis of basal cell carcinoma, the most common human cancer worldwide. It was hypothesized that FOXM1 induces cancer initiation through stem/progenitor cell expansion. What's more, FOXM1 has been shown to modulate the epigenome. It was found that overexpression of FOXM1 brain washes" normal cells to adopt cancer-like epigenome."

Images



Anti- FOXM1 antibody, ABO11885, Western blottingAll lanes: Anti FOXM1 (ABO11885) at 0.5ug/mlLane 1: HELA Whole Cell Lysate at 40ugLane 2: COLO320 Whole Cell Lysate at 40ugLane 3: SW620 Whole Cell Lysate at 40ugLane 4: SKOV Whole Cell Lysate at 40ugLane 5: MCD-7 Whole Cell Lysate at 40ugPredicted bind size: 84KDObserved bind size: 84KD



Anti- FOXM1 antibody, ABO11885, Western blottingAll lanes: Anti FOXM1 (ABO11885) at 0.5ug/mlWB: Recombinant Human FOXM1 Protein 0.5ngPredicted bind size: 36KDObserved bind size: 36KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.