

Anti-GRIA1 Picoband Antibody

Catalog # ABO11895

Product Information

Application	WB, IHC-P
Primary Accession	P42261
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Glutamate receptor 1(GRIA1) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	2890
Other Names	Glutamate receptor 1, GluR-1, AMPA-selective glutamate receptor 1, GluR-A, GluR-K1, Glutamate receptor ionotropic, AMPA 1, GluA1, GRIA1, GLUH1, GLUR1
Calculated MW	101506
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat
Subcellular Localization	Cell membrane ; Multi-pass membrane protein . Endoplasmic reticulum membrane ; Multi-pass membrane protein . Cell junction, synapse, postsynaptic cell membrane ; Multi-pass membrane protein . Cell junction, synapse, postsynaptic cell membrane, postsynaptic density . Cell projection, dendrite . Cell projection, dendritic spine . Interaction with CACNG2, CNIH2 and CNIH3 promotes cell surface expression. .
Tissue Specificity	Widely expressed in brain.
Source	Eukaryota
Protein Name	Glutamate receptor 1
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	E.coli-derived human GRIA1 recombinant protein (Position: A19-R360). Human GRIA1 shares 98% amino acid (aa) sequence identity with both mouse and rat GRIA1.

Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.

Protein Information

Name	GRIA1 (HGNC:4571)
Function	Ionotropic glutamate receptor that functions as a ligand- gated cation channel, gated by L-glutamate and glutamatergic agonists such as alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA), quisqualic acid, and kainic acid (PubMed:1311100, PubMed:20805473, PubMed:21172611, PubMed:28628100, PubMed:35675825). L- glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse upon entry of monovalent and divalent cations such as sodium and calcium. The receptor then desensitizes rapidly and enters in a transient inactive state, characterized by the presence of bound agonist (By similarity). In the presence of CACNG2 or CACNG4 or CACNG7 or CACNG8, shows resensitization which is characterized by a delayed accumulation of current flux upon continued application of L- glutamate (PubMed:21172611). Resensitization is blocked by CNIH2 through interaction with CACNG8 in the CACNG8-containing AMPA receptors complex (PubMed:21172611). Calcium (Ca(2+)) permeability depends on subunits composition and, heteromeric channels containing edited GRIA2 subunit are calcium-impermeable. Also permeable to other divalents cations such as strontium(2+) and magnesium(2+) and monovalent cations such as potassium(1+) and lithium(1+) (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein. Endoplasmic reticulum membrane {ECO:0000250 UniProtKB:P19490}; Multi-pass membrane protein {ECO:0000250 UniProtKB:P19490}. Postsynaptic cell membrane; Multi-pass membrane protein. Postsynaptic density membrane {ECO:0000250 UniProtKB:P23818}; Multi-pass membrane protein {ECO:0000250 UniProtKB:P23818}. Cell projection, dendrite {ECO:0000250 UniProtKB:P23818}. Cell projection, dendritic spine {ECO:0000250 UniProtKB:P23818}. Early endosome membrane {ECO:0000250 UniProtKB:P19490}; Multi-pass membrane protein {ECO:0000250 UniProtKB:P19490}. Recycling endosome membrane {ECO:0000250 UniProtKB:P19490}; Multi-pass membrane protein {ECO:0000250 UniProtKB:P19490}. Presynapse {ECO:0000250 UniProtKB:P23818}. Synapse {ECO:0000250 UniProtKB:P23818} Note=Interaction with CACNG2, CNIH2 and CNIH3 promotes cell surface expression. Colocalizes with PDLIM4 in early endosomes. Displays a somatodendritic localization and is excluded from axons in neurons (By similarity). Localized to cone photoreceptor pedicles (By similarity) {ECO:0000250 UniProtKB:P19490, ECO:0000250 UniProtKB:P23818}
Tissue Location	Widely expressed in brain.

Background

GLUR1, Glutamate receptor 1, is a protein that in humans is encoded by the GLUR1 gene. GLUR1 mRNA is widely expressed in human brain. Glutamate receptors are the predominant excitatory neurotransmitter receptors in the mammalian brain and are activated in a variety of normal neurophysiologic processes. The classification of glutamate receptors is based on their activation by different pharmacologic agonists. The GRIA1 belongs to a family of alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate (AMPA) receptors. Each of the members (GRIA1-4) include flip and flop isoforms generated by alternative RNA splicing. The receptor subunits encoded by each isoform vary in their signal transduction properties. The isoform presented here is the flop isoform. In situ hybridization experiments showed that human GRIA1 mRNA is present in granule and pyramidal cells in the hippocampal formation.


Images




Anti- GRIA1 antibody, ABO11895, Western blottingAll lanes: Anti GRIA1 (ABO11895) at 0.5ug/mlWB: Recombinant Human GRIA1 Protein 0.5ngPredicted bind size: 40KDObserved bind size: 40KD




Anti- GRIA1 antibody, ABO11895, Western blottingAll lanes: Anti GRIA1 (ABO11895) at 0.5ug/mlLane 1: Rat Brain Tissue Lysate at 50ugLane 2: Mouse Brain Tissue Lysate at 50ugPredicted bind size: 101KDObserved bind size: 101KD



Anti- GRIA1 antibody, ABO11895, IHC(P)IHC(P): Rat Brain Tissue



Anti- GRIA1 antibody, ABO11895, IHC(P)IHC(P): Mouse Brain Tissue



Anti- GRIA1 antibody, ABO11895, IHC(P)IHC(P): Human Meningeoma Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.