

Anti-FAS Antibody

Catalog # ABO11943

Product Information

Application	WB, IHC-P
Primary Accession	P25445
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Tumor necrosis factor receptor superfamily member 6(FAS) detection. Tested with WB, IHC-P in Human.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	355
Other Names	Tumor necrosis factor receptor superfamily member 6, Apo-1 antigen, Apoptosis-mediating surface antigen FAS, FASLG receptor, CD95, FAS, APT1, FAS1, TNFRSF6
Calculated MW	37732
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat Western blot, 0.1-0.5 µg/ml, Human
Subcellular Localization	Isoform 1: Cell membrane; Single-pass type I membrane protein.
Tissue Specificity	Isoform 1 and isoform 6 are expressed at equal levels in resting peripheral blood mononuclear cells. After activation there is an increase in isoform 1 and decrease in the levels of isoform 6. .
Source	Eukaryota
Protein Name	Tumor necrosis factor receptor superfamily member 6
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E.coli-derived human Fas recombinant protein (Position: Q26-N173). Human Fas shares 55% and 59% amino acid (aa) sequences identity with mouse and rat Fas, respectively.
Purification	Immunogen affinity purified.

Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.
Sequence Similarities	Contains 1 death domain.

Protein Information

Name	FAS
Synonyms	APT1, FAS1, TNFRSF6
Function	Receptor for TNFSF6/FASLG. The adapter molecule FADD recruits caspase CASP8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs CASP8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. FAS-mediated apoptosis may have a role in the induction of peripheral tolerance, in the antigen-stimulated suicide of mature T-cells, or both. The secreted isoforms 2 to 6 block apoptosis (in vitro).
Cellular Location	[Isoform 1]: Cell membrane; Single-pass type I membrane protein. Membrane raft [Isoform 3]: Secreted. [Isoform 5]: Secreted.
Tissue Location	Isoform 1 and isoform 6 are expressed at equal levels in resting peripheral blood mononuclear cells. After activation there is an increase in isoform 1 and decrease in the levels of isoform 6.

Background

FAS(also known as surface antigen APO1 or CD95) is a member of the tumour-necrosis receptor factor family of death receptors. It acts as an inducer of both neurite growth in vitro and accelerated recovery after nerve injury in vivo. FAS antigen is expressed and functional on papillary thyroid cancer cells and this may have potential therapeutic significance. The FAS antigen shows structural homology with a number of cell surface receptors, including tumor necrosis factor(TNF) receptors and the low-affinity nerve growth factor receptor(NGFR) and it is mapped to 10q24.1. The FAS and FASL system plays a key role in regulating apoptotic cell death and corruption of this signalling pathway has been shown to participate in immune escape and tumorigenesis.

Images




Anti- FAS Picoband antibody, ABO11943, Western blotting
All lanes: Anti FAS (ABO11943) at 0.5ug/ml
WB: Recombinant Human FAS Protein 0.5ng
Predicted bind size: 35KD
Observed bind size: 35KD

Anti- FAS Picoband antibody, ABO11943, Western



blotting All lanes: Anti FAS (ABO11943) at 0.5ug/ml
 Lane 1: HELA Whole Cell Lysate at 40ug
 Lane 2: JURKAT Whole Cell Lysate at 40ug
 Lane 3: A549 Whole Cell Lysate at 40ug
 Lane 4: SMMC Whole Cell Lysate at 40ug
 Lane 5: K562 Whole Cell Lysate at 40ug
 Predicted bind size: 37KD
 Observed bind size: 37KD



Anti- FAS Picoband antibody, ABO11943, IHC(P)
 Human Lung Cancer Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.