

Anti-ABCB4 Picoband Antibody

Catalog # ABO11966

Product Information

Application	WB, IHC-P
Primary Accession	P21439
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Phosphatidylcholine translocator ABCB4(ABCB4) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	5244
Other Names	Phosphatidylcholine translocator ABCB4, ATP-binding cassette sub-family B member 4 {ECO:0000312 HGNC:HGNC:45}, Multidrug resistance protein 3, 3.6.3.44, P-glycoprotein 3, ABCB4 (HGNC:45)
Calculated MW	141523
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5 µg/ml, Human
Subcellular Localization	Cell membrane; Multi-pass membrane protein.
Source	Eukaryota
Protein Name	Phosphatidylcholine translocator ABCB4
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E.coli-derived human ABCB4 recombinant protein (Position: A601-A720). Human ABCB4 shares 79% amino acid (aa) sequence identity with both mouse and rat ABCB4.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also

Sequence Similarities

be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.

Belongs to the ABC transporter superfamily, ABCB family, Multidrug resistance exporter (TC 3.A.1.201) subfamily.

Protein Information

Name ABCB4 ([HGNC:45](#))


Function [Isoform 1]: Energy-dependent phospholipid efflux translocator that acts as a positive regulator of biliary lipid secretion. Functions as a floppase that translocates specifically phosphatidylcholine (PC) from the inner to the outer leaflet of the canalicular membrane bilayer into the canaliculi of hepatocytes. Translocation of PC makes the biliary phospholipids available for extraction into the canaliculi lumen by bile salt mixed micelles and therefore protects the biliary tree from the detergent activity of bile salts (PubMed:[17523162](#), PubMed:[21820390](#), PubMed:[23468132](#), PubMed:[24594635](#), PubMed:[24723470](#), PubMed:[24806754](#), PubMed:[31873305](#), PubMed:[7957936](#), PubMed:[8898203](#), PubMed:[9366571](#)). Plays a role in the recruitment of phosphatidylcholine (PC), phosphatidylethanolamine (PE) and sphingomyelin (SM) molecules to nonraft membranes and to further enrichment of SM and cholesterol in raft membranes in hepatocytes (PubMed:[23468132](#)). Required for proper phospholipid bile formation (By similarity). Indirectly involved in cholesterol efflux activity from hepatocytes into the canalicular lumen in the presence of bile salts in an ATP-dependent manner (PubMed:[24045840](#)). Promotes biliary phospholipid secretion as canaliculi-containing vesicles from the canalicular plasma membrane (PubMed:[28012258](#), PubMed:[9366571](#)). In cooperation with ATP8B1, functions to protect hepatocytes from the deleterious detergent activity of bile salts (PubMed:[21820390](#)). Does not confer multidrug resistance (By similarity).

Cellular Location Cell membrane; Multi-pass membrane protein {ECO:0000255|PROSITE-ProRule:PRU00441}. Apical cell membrane; Multi-pass membrane protein {ECO:0000255|PROSITE-ProRule:PRU00441}. Membrane raft. Cytoplasm Cytoplasmic vesicle, clathrin-coated vesicle {ECO:0000250|UniProtKB:Q08201}. Note=Localized at the apical canalicular membrane of the epithelial cells lining the lumen of the bile canaliculi and biliary ductules (By similarity). Transported from the Golgi to the apical bile canalicular membrane in a RACK1-dependent manner (PubMed:19674157). Redistributed into pseudocanaliculi formed between cells in a bezafibrate- or PPARA-dependent manner (PubMed:15258199). Localized preferentially in lipid nonraft domains of canalicular plasma membranes (PubMed:23468132) {ECO:0000250|UniProtKB:P21440, ECO:0000269|PubMed:15258199, ECO:0000269|PubMed:19674157, ECO:0000269|PubMed:23468132}


Background

Adenosine triphosphate-binding cassette subfamily B, member 4 (ABCB4), also called MDR3, is a protein that in humans is encoded by the ABCB4 gene. The membrane-associated protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. The ABCB4 gene contains 28 exons and 27 of these contain coding sequences for the two homologous halves of the protein that correlate with functional domains. ABCB4 gene mutations represent a genetic factor involved in this peculiar form of cholesterol gallstone disease in adults.


Images




Anti- ABCB4 Picoband antibody, ABO11966, Western blotting
 All lanes: Anti ABCB4 (ABO11966) at 0.5ug/ml
 Lane 1: MCF-7 Whole Cell Lysate at 40ug
 Lane 2: SW620 Whole Cell Lysate at 40ug
 Lane 3: 22RV1 Whole Cell Lysate at 40ug
 Lane 4: SKOV Whole Cell Lysate at 40ug
 Predicted bind size: 142KD
 Observed bind size: 142KD



Anti- ABCB4 Picoband antibody, ABO11966, IHC(P)
 Mouse Intestine Tissue



Anti- ABCB4 Picoband antibody, ABO11966, IHC(P)
 Rat Liver Tissue



Anti- ABCB4 Picoband antibody, ABO11966, IHC(P)
 Human Mammary Cancer Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.