

Anti-NUMB Picoband Antibody

Catalog # ABO11992

Product Information

Application	WB
Primary Accession	P49757
Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Protein numb homolog(NUMB) detection. Tested with WB in Human;Mouse.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	8650
Other Names	Protein numb homolog, h-Numb, Protein S171, NUMB
Calculated MW	70804
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Mouse
Subcellular Localization	Membrane; Peripheral membrane protein.
Source	Eukaryota
Protein Name	Protein numb homolog
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human NUMB(194-220aa SFRVTTATEQAEREEIMKQMQDAKKAE), different from the related mouse and rat sequences by one amino acid.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.
Sequence Similarities	Contains 1 PID domain.

Protein Information

Name	NUMB (HGNC:8060)
Function	Regulates clathrin-mediated receptor endocytosis (PubMed: 18657069). Plays a role in the process of neurogenesis (By similarity). Required throughout embryonic neurogenesis to maintain neural progenitor cells, also called radial glial cells (RGCs), by allowing their daughter cells to choose progenitor over neuronal cell fate (By similarity). Not required for the proliferation of neural progenitor cells before the onset of neurogenesis. Also involved postnatally in the subventricular zone (SVZ) neurogenesis by regulating SVZ neuroblasts survival and ependymal wall integrity (By similarity). May also mediate local repair of brain ventricular wall damage (By similarity).
Cellular Location	Cell membrane; Peripheral membrane protein; Cytoplasmic side. Endosome membrane; Peripheral membrane protein; Cytoplasmic side. Note=Localizes to perinuclear endosomes in an AAK1-dependent manner.

Background

Protein numb homolog is a protein that in humans is encoded by the NUMB gene. It is mapped to 14q24.2 to q24.3. The encoded protein, whose degradation is induced in a proteasome-dependent manner by MDM2, is a membrane-bound protein that has been shown to associate with EPS15, LNX1, and NOTCH1. The primary function of Numb in cell differentiation is as an inhibitor of Notch signaling which is essential for maintaining self-renewal potential in stem and progenitor cells. Numb also plays a crucial role in asymmetrical cell division during development, allowing for differential cell fate specification in the central and peripheral nervous systems. What is more, the numb gene protein product controls binary cell fate decisions in the peripheral and central nervous systems of both invertebrates and mammals during neurogenesis.

Images



Anti- NUMB Picoband antibody, ABO11992, Western blotting
All lanes: Anti NUMB (ABO11992) at 0.5ug/ml
Lane 1: U87 Whole Cell Lysate at 40ug
Lane 2: NEURO Whole Cell Lysate at 40ug
Predicted bind size: 71KD
Observed bind size: 71KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.