

Anti-NMDAR2A Picoband Antibody

Catalog # ABO12026

Product Information

Application	WB
Primary Accession	Q12879
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Glutamate receptor ionotropic, NMDA 2A (GRIN2A) detection. Tested with WB in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	2903
Other Names	Glutamate receptor ionotropic, NMDA 2A, GluN2A, Glutamate [NMDA] receptor subunit epsilon-1, N-methyl D-aspartate receptor subtype 2A, NMDAR2A, NR2A, hNR2A, GRIN2A, NMDAR2A
Calculated MW	165283
Application Details	Western blot, 0.1-0.5 µg/ml, Mouse, Rat, Human
Subcellular Localization	Cell membrane; Multi-pass membrane protein. Cell junction, synapse, postsynaptic cell membrane; Multi-pass membrane protein.
Source	Eukaryota
Protein Name	Glutamate receptor ionotropic, NMDA 2A
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E.coli-derived human NMDAR2A recombinant protein (Position: D958-R1300). Human NMDAR2A shares 89% and 90% amino acid (aa) sequence identity with mouse and rat NMDAR2A, respectively.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.

Protein Information

Name	GRIN2A (HGNC:4585)
Synonyms	NMDAR2A
Function	Component of N-methyl-D-aspartate (NMDA) receptors (NMDARs) that function as heterotetrameric, ligand-gated cation channels with high calcium permeability and voltage-dependent block by Mg(2+) (PubMed:20890276, PubMed:23933818, PubMed:23933819, PubMed:23933820, PubMed:24504326, PubMed:26875626, PubMed:26919761, PubMed:28242877, PubMed:36117210, PubMed:38538865, PubMed:8768735). NMDARs participate in synaptic plasticity for learning and memory formation by contributing to the slow phase of excitatory postsynaptic current, long-term synaptic potentiation, and learning (By similarity). Channel activation requires binding of the neurotransmitter L-glutamate to the GluN2 subunit, glycine or D-serine binding to the GluN1 subunit, plus membrane depolarization to eliminate channel inhibition by Mg(2+) (PubMed:23933818, PubMed:23933819, PubMed:23933820, PubMed:24504326, PubMed:26875626, PubMed:26919761, PubMed:27288002, PubMed:28095420, PubMed:28105280, PubMed:28126851, PubMed:28182669, PubMed:29644724, PubMed:38307912, PubMed:8768735). NMDARs mediate simultaneously the potassium efflux and the influx of calcium and sodium (By similarity). Each GluN2 subunit confers differential attributes to channel properties, including activation, deactivation and desensitization kinetics, pH sensitivity, Ca2(+) permeability, and binding to allosteric modulators (PubMed:26875626, PubMed:26919761). Participates in the synaptic plasticity regulation through activation by the L- glutamate released by BEST1, into the synaptic cleft, upon F2R/PAR-1 activation in astrocyte (By similarity).
Cellular Location	Cell projection, dendritic spine {ECO:0000250 UniProtKB:Q00959}. Cell membrane; Multi-pass membrane protein. Synapse {ECO:0000250 UniProtKB:P35436} Postsynaptic cell membrane {ECO:0000250 UniProtKB:Q00959}; Multi-pass membrane protein. Cytoplasmic vesicle membrane {ECO:0000250 UniProtKB:P35436}. Note=Expression at the dendrite cell membrane and at synapses is regulated by SORCS2 and the retromer complex. {ECO:0000250 UniProtKB:P35436}

Background

GRIN2A is also known as N-methyl-D-aspartate receptor channel, subunit epsilon-1(NMDAR2A). This gene encodes a member of the glutamate-gated ion channel protein family. The encoded protein is an N-methyl-D-aspartate (NMDA) receptor subunit. NMDA receptors are both ligand-gated and voltage-dependent, and are involved in long-term potentiation, an activity-dependent increase in the efficiency of synaptic transmission thought to underlie certain kinds of memory and learning. These receptors are permeable to calcium ions, and activation results in a calcium influx into post-synaptic cells, which results in the activation of several signaling cascades. Disruption of this gene is associated with focal epilepsy and speech disorder with or without mental retardation. Alternative splicing results in multiple transcript variants.

Images

Anti- NMDAR2A Picoband antibody, ABO12026, Western blottingAll lanes: Anti NMDAR2A (ABO12026) at



0.5ug/ml
Lane 1: Rat Brain Tissue Lysate at 50ug
Lane 2: Mouse Brain Tissue Lysate at 50ug
Predicted bind size: 165KD
Observed bind size: 165KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.