

Anti-Caspase-2 Picoband Antibody

Catalog # ABO12059

Product Information

Application	WB
Primary Accession	P42575
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Caspase-2(CASP2) detection. Tested with WB in Human;Mouse;Rat.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	835
Other Names	Caspase-2, CASP-2, 3.4.22.55, Neural precursor cell expressed developmentally down-regulated protein 2, NEDD-2, Protease ICH-1, Caspase-2 subunit p18, Caspase-2 subunit p13, Caspase-2 subunit p12, CASP2, ICH1, NEDD2
Calculated MW	50685
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat
Tissue Specificity	Expressed at higher levels in the embryonic lung, liver and kidney than in the heart and brain. In adults, higher level expression is seen in the placenta, lung, kidney, and pancreas than in the heart, brain, liver and skeletal muscle.
Source	Eukaryota
Protein Name	Caspase-2
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human CASP2(378-409aa RNTKRGSWYIEALAQVFSEACDMHVADMLVK), different from the related mouse and rat sequences by one amino acid.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins.
Storage	At -20° C for one year. After reconstitution, at 4° C for one month. It can also

Sequence Similarities

be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
Belongs to the peptidase C14A family.


Protein Information

Name	CASP2
Synonyms	ICH1, NEDD2
Function	Is a regulator of the cascade of caspases responsible for apoptosis execution (PubMed: 11156409 , PubMed: 15073321 , PubMed: 8087842). Might function by either activating some proteins required for cell death or inactivating proteins necessary for cell survival (PubMed: 15073321). Associates with PIDD1 and CRADD to form the PIDDosome, a complex that activates CASP2 and triggers apoptosis in response to genotoxic stress (PubMed: 15073321).
Tissue Location	Expressed at higher levels in the embryonic lung, liver and kidney than in the heart and brain. In adults, higher level expression is seen in the placenta, lung, kidney, and pancreas than in the heart, brain, liver and skeletal muscle

Background

CASP2 is equal to Caspase-2. And Caspase-2, which is involved in stress-induced apoptosis, is recruited into a large protein complex, the molecular composition of which remains elusive. It is showed that activation of caspase-2 occurs in a complex that contains the death domain-containing protein PIDD, whose expression is induced by p53, and the adaptor protein RAIDD. Increased PIDD expression resulted in spontaneous activation of caspase-2 and sensitization to apoptosis by genotoxic stimuli. Caspase-2 acts both as a positive and negative cell death effector, depending upon cell lineage and stage of development.

Images



Anti- Caspase-2 Picoband antibody, ABO12059, Western blottingAll lanes: Anti Caspase-2 (ABO12059) at 0.5ug/mlLane 1: Rat Lung Tissue Lysate at 50ugLane 2: Mouse Liver Tissue Lysate at 50ugLane 3: A549 Whole Cell Lysate at 40ugLane 4: PANC Whole Cell Lysate at 40ugLane 5: 293T Whole Cell Lysate at 40ugPredicted bind size: 18KDObserved bind size: 18KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.