

Anti-STIM1 Picoband Antibody

Catalog # ABO12097

Product Information

Application	WB, IHC-P
Primary Accession	Q13586
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Stromal interaction molecule 1(STIM1) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	6786
Other Names	Stromal interaction molecule 1, STIM1, GOK {ECO:0000303 PubMed:9377559}
Calculated MW	77423
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5 μ g/ml, Human, Mouse, Rat
Subcellular Localization	Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein. Cytoplasm, cytoskeleton. Translocates from the endoplasmic reticulum to the cell membrane in response to a depletion of intracellular calcium and is detected at punctae corresponding to junctions between the endoplasmic reticulum and the cell membrane. Associated with the microtubule network at the growing distal tip of microtubules.
Tissue Specificity	Ubiquitously expressed in various human primary cells and tumor cell lines. .
Source	Eukaryota
Protein Name	Stromal interaction molecule 1
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human STIM1(45-74aa AAEFCRIDKPLCHSEDEKLSFEAVRNIIHKL), identical to the related mouse and rat sequences.

Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins.
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Contains 1 EF-hand domain.

Protein Information

Name	STIM1
Synonyms	GOK {ECO:0000303 PubMed:9377559}
Function	Acts as a Ca(2+) sensor that gates two major inward rectifying Ca(2+) channels at the plasma membrane: Ca(2+) release- activated Ca(2+) (CRAC) channels and arachidonate-regulated Ca(2+)- selective (ARC) channels (PubMed: 15866891 , PubMed: 16005298 , PubMed: 16208375 , PubMed: 16537481 , PubMed: 16733527 , PubMed: 16766533 , PubMed: 16807233 , PubMed: 18854159 , PubMed: 19182790 , PubMed: 19249086 , PubMed: 19622606 , PubMed: 19706554 , PubMed: 22464749 , PubMed: 24069340 , PubMed: 24351972 , PubMed: 24591628 , PubMed: 25326555 , PubMed: 26322679 , PubMed: 28219928 , PubMed: 32415068). Plays a role in mediating store-operated Ca(2+) entry (SOCE), a Ca(2+) influx following depletion of intracellular Ca(2+) stores. Upon Ca(2+) depletion, translocates from the endoplasmic reticulum to the plasma membrane where it activates CRAC channel pore-forming subunits ORA1, ORA2 and ORAI3 to generate sustained and oscillatory Ca(2+) entry (PubMed: 16208375 , PubMed: 16537481 , PubMed: 32415068). Involved in enamel formation (PubMed: 24621671). Plays a role in regulating the type I interferon response by retaining the signaling adapter STING at the endoplasmic reticulum (PubMed: 30643259).
Cellular Location	Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein. Cytoplasm, cytoskeleton. Sarcoplasmic reticulum. Note=Translocates from the endoplasmic reticulum to the cell membrane in response to a depletion of intracellular calcium and is detected at punctae corresponding to junctions between the endoplasmic reticulum and the cell membrane (PubMed:16005298, PubMed:16208375, PubMed:18854159, PubMed:19182790, PubMed:19249086). Associated with the microtubule network at the growing distal tip of microtubules (PubMed:19632184). Colocalizes with ORAI1 at the cell membrane (PubMed:27185316). Colocalizes preferentially with CASQ1 at endoplasmic reticulum in response to a depletion of intracellular calcium (PubMed:27185316)
Tissue Location	Ubiquitously expressed in various human primary cells and tumor cell lines.

Background


Stromal interaction molecule 1 is a protein that in humans is encoded by the STIM1 gene. STIM1 has a single transmembranedomain, and is localized to the endoplasmic reticulum, and to a lesser extent to the plasma membrane. This gene encodes a type 1 transmembrane protein that mediates Ca²⁺ influx after depletion of intracellular Ca²⁺ stores by gating of store-operated Ca²⁺ influx channels (SOCs). It is one of several genes located in the imprinted gene domain of 11p15.5, an important tumor-suppressor gene region. Alterations in this region have been associated with the Beckwith-Wiedemann syndrome, Wilms tumor,

rhabdomyosarcoma, adrenocortical carcinoma, and lung, ovarian, and breast cancer. This gene may play a role in malignancies and disease that involve this region, as well as early hematopoiesis, by mediating attachment to stromal cells. Mutations in this gene are associated with fatal classic Kaposi sarcoma, immunodeficiency due to defects in store-operated calcium entry (SOCE) in fibroblasts, ectodermal dysplasia and tubular aggregate myopathy. This gene is oriented in a head-to-tail configuration with the ribonucleotide reductase 1 gene (RRM1), with the 3' end of this gene situated 1.6 kb from the 5' end of the RRM1 gene. Alternative splicing of this gene results in multiple transcript variants.


Images




Anti- STIM1 Picoband antibody, ABO12097, Western blotting
All lanes: Anti STIM1 (ABO12097) at 0.5ug/ml
Lane 1: Rat Liver Tissue Lysate at 50ug
Lane 2: Mouse Liver Tissue Lysate at 50ug
Lane 3: Human Placenta Tissue Lysate at 50ug
Lane 4: HELA Whole Cell Lysate at 40ug
Lane 5: SMMC Whole Cell Lysate at 40ug
Lane 6: HEPG2 Whole Cell Lysate at 40ug
Predicted bind size: 77KD
Observed bind size: 100KD



Anti- STIM1 Picoband antibody, ABO12097, IHC(P)
IHC(P): Mouse Intestine Tissue



Anti- STIM1 Picoband antibody, ABO12097, IHC(P)
IHC(P): Rat Intestine Tissue



Anti- STIM1 Picoband antibody, ABO12097, IHC(P)
IHC(P): Human Mammary Cancer Tissue