

Anti-APH1a Picoband Antibody

Catalog # ABO12111

Product Information

Application	WB
Primary Accession	Q96BI3
Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Gamma-secretase subunit APH-1A(APH1A) detection. Tested with WB in Human;Mouse.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	51107
Other Names	Gamma-secretase subunit APH-1A, APH-1a, Aph-1alpha, Presenilin-stabilization factor, APH1A, PSF
Calculated MW	28996
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Mouse
Subcellular Localization	Endoplasmic reticulum membrane; Multi-pass membrane protein. Golgi apparatus, Golgi stack membrane; Multi- pass membrane protein. Predominantly located in the endoplasmic reticulum and in the cis-Golgi.
Tissue Specificity	Widely expressed. Expressed in leukocytes, lung, placenta, small intestine, liver, kidney, spleen thymus, skeletal muscle, heart and brain. Isoform 1 and isoform 2 are nearly expressed at the same level. .
Source	Eukaryota
Protein Name	Gamma-secretase subunit APH-1A
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Na ₃ N.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human APH1a (236-265aa LRSIQRSLLCRRQEDSRVMVYSALRIPPED), different from the related mouse sequence by one amino acid.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins

Storage	At -20 °C for one year. After r° Constitution, at 4 °C for one month. It° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the APH-1 family.

Protein Information


Name	APH1A
Synonyms	PSF
Function	Non-catalytic subunit of the gamma-secretase complex, an endoprotease complex that catalyzes the intramembrane cleavage of integral membrane proteins such as Notch receptors and APP (amyloid- beta precursor protein) (PubMed: 12297508 , PubMed: 12522139 , PubMed: 12679784 , PubMed: 12763021 , PubMed: 25043039 , PubMed: 26280335 , PubMed: 30598546 , PubMed: 30630874). Required for normal gamma-secretase assembly (PubMed: 12471034 , PubMed: 12522139 , PubMed: 12763021 , PubMed: 19369254). The gamma-secretase complex plays a role in Notch and Wnt signaling cascades and regulation of downstream processes via its role in processing key regulatory proteins, and by regulating cytosolic CTNNB1 levels (Probable).
Cellular Location	Endoplasmic reticulum membrane; Multi-pass membrane protein. Golgi apparatus, Golgi stack membrane; Multi-pass membrane protein. Note=Predominantly located in the endoplasmic reticulum and in the cis-Golgi
Tissue Location	Widely expressed. Expressed in leukocytes, lung, placenta, small intestine, liver, kidney, spleen thymus, skeletal muscle, heart and brain. Isoform 1 and isoform 2 are nearly expressed at the same level.

Background

APH1a encodes a component of the gamma secretase complex that cleaves integral membrane proteins such as Notch receptors and beta-amyloid precursor protein. The gamma secretase complex contains this gene product, or the paralogous anterior pharynx defective 1 homolog B (APH1B), along with the presenilin, nicastrin, and presenilin enhancer-2 proteins. The precise function of this seven-transmembrane-domain protein is unknown though it is suspected of facilitating the association of nicastrin and presenilin in the gamma secretase complex as well as interacting with substrates of the gamma secretase complex prior to their proteolytic processing. Polymorphisms in a promoter region of this gene have been associated with an increased risk for developing sporadic Alzheimer's disease. Alternative splicing results in multiple protein-coding and non-protein-coding transcript variants.

Images

Anti- APH1a Picoband antibody, ABO12111, Western blottingAll lanes: Anti APH1a (ABO12111) at 0.5ug/mlLane 1: Mouse Lung Tissue Lysate at 50ugLane 2: Mouse Liver Tissue Lysate at 50ugLane 3: SW620 Whole Cell Lysate at 40ugLane 4: SMMC Whole Cell Lysate at 40ugLane 5: Human Placenta Tissue Lysate at 50ugPredicted bind size: 29KDObserved bind size: 29KD



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.