

Anti-Band 3 Picoband Antibody

Catalog # ABO12127

Product Information

Application	WB, IHC-P, IHC-F
Primary Accession	P02730
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Band 3 anion transport protein(SLC4A1) detection. Tested with WB, IHC-P, IHC-F in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	6521
Other Names	Band 3 anion transport protein, Anion exchange protein 1, AE 1, Anion exchanger 1, Solute carrier family 4 member 1, CD233, SLC4A1, AE1, DI, EPB3
Calculated MW	101792
Application Details	Immunohistochemistry(Frozen Section), 0.5-1 μ g/ml, Human, - Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5 μ g/ml, Human, Rat
Subcellular Localization	Cell membrane; Multi-pass membrane protein. Basolateral cell membrane; Multi-pass membrane protein. Detected in the erythrocyte cell membrane and on the basolateral membrane of alpha-intercalated cells in the collecting duct in the kidney.
Tissue Specificity	Detected in erythrocytes (at protein level). Erythrocytes. .
Source	Eukaryota
Protein Name	Band 3 anion transport protein
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E. coli-derived human Band 3 (Position: E28-N365). Human Band 3 shares 75.7% and 74.5% amino acid (aa) sequence identity with mouse and rat Band 3, respectively.
Purification	Immunogen affinity purified.

Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the anion exchanger (TC 2.A.31) family.

Protein Information


Name	SLC4A1 (HGNC:11027)
Synonyms	AE1, DI, EPB3
Function	Functions both as a transporter that mediates electroneutral anion exchange across the cell membrane and as a structural protein (PubMed: 10926824 , PubMed: 14734552 , PubMed: 1538405 , PubMed: 16227998 , PubMed: 20151848 , PubMed: 24121512 , PubMed: 28387307 , PubMed: 35835865). Component of the ankyrin-1 complex of the erythrocyte membrane; required for normal flexibility and stability of the erythrocyte membrane and for normal erythrocyte shape via the interactions of its cytoplasmic domain with cytoskeletal proteins, glycolytic enzymes, and hemoglobin (PubMed: 1538405 , PubMed: 20151848 , PubMed: 35835865). Functions as a transporter that mediates the 1:1 exchange of inorganic anions across the erythrocyte membrane. Mediates chloride-bicarbonate exchange in the kidney, and is required for normal acidification of the urine (PubMed: 10926824 , PubMed: 14734552 , PubMed: 16227998 , PubMed: 24121512 , PubMed: 28387307).
Cellular Location	Cell membrane; Multi-pass membrane protein. Basolateral cell membrane; Multi-pass membrane protein. Note=Detected in the erythrocyte cell membrane and on the basolateral membrane of alpha-intercalated cells in the collecting duct in the kidney.
Tissue Location	Detected in erythrocytes (at protein level).

Background


Band 3 is also known as SLC4A1. The protein encoded by this gene is part of the anion exchanger (AE) family and is expressed in the erythrocyte plasma membrane, where it functions as a chloride/bicarbonate exchanger involved in carbon dioxide transport from tissues to lungs. The protein comprises two domains that are structurally and functionally distinct. The N-terminal 40kDa domain is located in the cytoplasm and acts as an attachment site for the red cell skeleton by binding ankyrin. The glycosylated C-terminal membrane-associated domain contains 12-14 membrane spanning segments and carries out the stilbene disulphonate-sensitive exchange transport of anions. The cytoplasmic tail at the extreme C-terminus of the membrane domain binds carbonic anhydrase II. The encoded protein associates with the red cell membrane protein glycophorin A and this association promotes the correct folding and translocation of the exchanger. This protein is predominantly dimeric but forms tetramers in the presence of ankyrin. Many mutations in this gene are known in man, and these mutations can lead to two types of disease: destabilization of red cell membrane leading to hereditary spherocytosis, and defective kidney acid secretion leading to distal renal tubular acidosis. Other mutations that do not give rise to disease result in novel blood group antigens, which form the Diego blood group system.

Images


Anti- Band 3 Picoband antibody, ABO12127, Western




blottingAll lanes: Anti Band 3 (ABO12127) at 0.5ug/ml
 Lane 1: Rat Brain Tissue Lysate at 50ug
 Lane 2: Rat Kidney Tissue Lysate at 50ug
 Lane 3: Rat Liver Tissue Lysate at 50ug
 Lane 4: Rat Spleen Tissue Lysate at 50ug
 Lane 5: Human Placenta Tissue Lysate at 50ug
 Predicted bind size: 102K
 Observed bind size: 102KD




Anti- Band 3 Picoband antibody, ABO12127,IHC(P)
 Rat Kidney Tissue



Anti- Band 3 Picoband antibody, ABO12127,IHC(P)
 Human Placenta Tissue



Anti- Band 3 Picoband antibody, ABO12127,IHC(P)
 Mouse Kidney Tissue



Anti- Band 3 Picoband antibody, ABO12127,IHC(F)
 Human Placenta Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.