10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-TRAF2 Picoband Antibody

Catalog # ABO12204

Product Information

Application	WB, IHC-P
Primary Accession	Q12933
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for TNF receptor-associated factor 2(TRAF2) detection. Tested with WB, IHC-P in Human.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	7186
Other Names	TNF receptor-associated factor 2, 6.3.2.-, E3 ubiquitin-protein ligase TRAF2, Tumor necrosis factor type 2 receptor-associated protein 3, TRAF2, TRAP3
Calculated MW	55859
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 □g/ml, Human, By Heat Western blot, 0.1-0.5 □g/ml, Human
Subcellular Localization	Cytoplasm .
Source	Eukaryota
Protein Name	TNF receptor-associated factor 2
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human TRAF2 (308-339aa RLDQDKIEALSSKVQQLERSIGLKDLAMADLE), different from the related mouse sequence by one amino acid.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20˚C for one year. After r˚Constitution, at 4˚C for one month. It˚Can also be aliquotted and stored frozen at -20˚C for a longer time.Avoid repeated

	freezing and thawing.
Sequence Similarities	Belongs to the TNF receptor-associated factor family. A subfamily.

Protein Information

Name	TRAF2 {ECO:0000303|PubMed:28489822, ECO:0000312|HGNC:HGNC:12032}
Function	E3 ubiquitin-protein ligase that regulates activation of NF- kappa-B and JNK and plays a central role in the regulation of cell survival and apoptosis (PubMed:10346818, PubMed:11784851, PubMed:12917689, PubMed:15383523, PubMed:18981220, PubMed:19150425, PubMed:19810754, PubMed:19918265, PubMed:19937093, PubMed:20047764, PubMed:20064526, PubMed:20385093, PubMed:20577214, PubMed:22212761). Catalyzes 'Lys-63'-linked ubiquitination of target proteins, such as BIRC3, IKBKE, MLST8, RIPK1 and TICAM1 (PubMed:23453969, PubMed:28489822). Is an essential constituent of several E3 ubiquitin- protein ligase complexes, where it promotes the ubiquitination of target proteins by bringing them into contact with other E3 ubiquitin ligases (PubMed:15383523, PubMed:18981220). Regulates BIRC2 and BIRC3 protein levels by inhibiting their autoubiquitination and subsequent degradation; this does not depend on the TRAF2 RING-type zinc finger domain (PubMed:11907583, PubMed:19506082). Plays a role in mediating activation of NF-kappa-B by EIF2AK2/PKR (PubMed:15121867). In complex with BIRC2 or BIRC3, promotes ubiquitination of IKBKE (PubMed:23453969). Acts as a regulator of mTORC1 and mTORC2 assembly by mediating 'Lys-63'-linked ubiquitination of MLST8, thereby inhibiting formation of the mTORC2 complex, while facilitating assembly of the mTORC1 complex (PubMed:28489822). Required for normal antibody isotype switching from IgM to IgG (By similarity).
Cellular Location	Cytoplasm


Background

TRAF2 (TNF Receptor-Associated Factor 2), also called TRAP, is a protein that in humans is encoded by the TRAF2 gene. The protein encoded by this gene is a member of the TNF receptor (TNFR) associated factor (TRAF) protein family. TRAF2 is a common signal transducer for TNFR2 and CD40 that mediates activation of NF-kappa-B. Rothe et al. (1996) identified ITRAF, which binds to TRAF1, TRAF2, and TRAF3, and that when overexpressed inhibits TRAF2-mediated NF-kappa-B activation. They proposed that ITRAF is an inhibitor of TRAF function that regulates TRAF protein activity by sequestering TRAFs in a latent state in the cytoplasm. Kanamori et al. (2002) found that mouse Traf2 interacted directly with T2bp, and they presented evidence that T2BP is involved in TNF-mediated signaling by its interaction with TRAF2.

Images

Anti- TRAF2 Picoband antibody, ABO12204, Western blottingAll lanes: Anti TRAF2 (ABO12204) at 0.5ug/mlLane 1: HELA Whole Cell Lysate at 40ugLane 2: JURKAT Whole Cell Lysate at 40ugPredicted bind size: 56KDObserved bind size: 56KD





Anti- TRAF2 Picoband antibody, ABO12204, IHC(P)IHC(P): Human Tonsil Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.