

Anti-IL-7 Antibody

Catalog # ABO12208

Product Information

Application	WB, E
Primary Accession	P13232
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Interleukin-7(IL7) detection. Tested with WB, ELISA in Human.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	3574
Other Names	Interleukin-7, IL-7, IL7
Calculated MW	20187
Application Details	Western blot, 0.1-0.5 µg/ml, Human, - ELISA , 0.1-0.5 µg/ml, Human
Subcellular Localization	Secreted.
Source	Eukaryota
Protein Name	Interleukin-7
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	E. coli-derived human IL7 recombinant protein (Position: D26-H177). Human IL7 shares 56.4% and 57% amino acid (aa) sequence identity with mouse and rat IL7, respectively.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20° C for one year. After reconstitution, at 4° C for one month. It can also be aliquotted and stored frozen at -20° C for a longer time. Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the IL-7/IL-9 family.


Protein Information

Name	IL7
Function	Hematopoietic cytokine that plays an essential role in the development, expansion, and survival of naive and memory T-cells and B- cells thereby regulating the number of mature lymphocytes and maintaining lymphoid homeostasis (PubMed: 25870237 , PubMed: 7527823). Mechanistically, exerts its biological effects through a receptor composed of IL7RA subunit and the cytokine receptor common subunit gamma/CSF2RG (PubMed: 8128231). Binding to the receptor leads to activation of various kinases including JAK1 or JAK3 depending on the cell type and subsequently propagation of signals through activation of several downstream signaling pathways including the PI3K/Akt/mTOR or the JAK-STAT5 (PubMed: 18523275 , PubMed: 20974963).
Cellular Location	Secreted.

Background

IL-7, Interleukin 7, is a protein that in humans is encoded by the IL-7 gene. IL-7 a hematopoietic growth factor secreted by stromal cells in the red marrow and thymus. It is also produced by keratinocytes, dendritic cells, hepatocytes, neurons, and epithelial cells but is not produced by lymphocytes. By the combination of approaches, IL-7 is located on 8q12-q13. IL-7 is critical for early T-cell development and homeostasis of naive and memory CD8-positive T cells. It signals through the IL-7 receptor complex, which consists of the IL7R-alpha chain and the common IL2R-gamma chain is critical for early T-cell development and homeostasis of naive and memory CD8 -positive T cells. It signals through the IL-7 receptor complex, which consists of the IL7R-alpha chain and the common IL2R-gamma chain.

Images



Anti- IL7 Picoband antibody, ABO12208, Western blottingAll lanes: Anti IL7 (ABO12208) at 0.5ug/mlWB: K562 Whole Cell Lysate at 40ugPredicted bind size: 20KDObserved bind size: 75KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.