

Anti-CYP1A1 Picoband Antibody

Catalog # ABO12231

Product Information

Application	WB, IHC-P, IHC-F
Primary Accession	P04798
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Cytochrome P450 1A1(CYP1A1) detection. Tested with WB, IHC-P, IHC-F in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	1543
Other Names	Cytochrome P450 1A1, 1.14.14.1, CYP1A1, Cytochrome P450 form 6, Cytochrome P450-C, Cytochrome P450-P1, CYP1A1
Calculated MW	58165
Application Details	Immunohistochemistry(Frozen Section), 0.5-1 μ g/ml, Human, Mouse, - Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5 μ g/ml, Human, Mouse, Rat
Subcellular Localization	Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein.
Tissue Specificity	Lung, lymphocytes and placenta.
Source	Eukaryota
Protein Name	Cytochrome P450 1A1
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Na ₃ N.
Immunogen	E.coli-derived human CYP1A1 recombinant protein (Position: H183-D320). Human CYP1A1 shares 81.2% amino acid (aa) sequence identity with both mouse and rat CYP1A1.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins

Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the cytochrome P450 family.

Protein Information


Name	CYP1A1 {ECO:0000303 PubMed:10681376, ECO:0000312 HGNC:HGNC:2595}
Function	A cytochrome P450 monooxygenase involved in the metabolism of various endogenous substrates, including fatty acids, steroid hormones and vitamins (PubMed:10681376, PubMed:11555828, PubMed:12865317, PubMed:14559847, PubMed:15041462, PubMed:15805301, PubMed:18577768, PubMed:19965576, PubMed:20972997). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed:10681376, PubMed:11555828, PubMed:12865317, PubMed:14559847, PubMed:15041462, PubMed:15805301, PubMed:18577768, PubMed:19965576, PubMed:20972997). Catalyzes the hydroxylation of carbon-hydrogen bonds. Exhibits high catalytic activity for the formation of hydroxysteroids from estrone (E1) and 17beta-estradiol (E2), namely 2-hydroxy E1 and E2, as well as D-ring hydroxylated E1 and E2 at the C15-alpha and C16- alpha positions (PubMed:11555828, PubMed:12865317, PubMed:14559847, PubMed:15805301). Displays different regioselectivities for polyunsaturated fatty acids (PUFA) hydroxylation (PubMed:15041462, PubMed:18577768). Catalyzes the epoxidation of double bonds of certain PUFA (PubMed:15041462, PubMed:19965576, PubMed:20972997). Converts arachidonic acid toward epoxyeicosatrienoic acid (EET) regioisomers, 8,9-, 11,12-, and 14,15-EET, that function as lipid mediators in the vascular system (PubMed:20972997). Displays an absolute stereoselectivity in the epoxidation of eicosapentaenoic acid (EPA) producing the 17(R),18(S) enantiomer (PubMed:15041462). May play an important role in all-trans retinoic acid biosynthesis in extrahepatic tissues. Catalyzes two successive oxidative transformation of all-trans retinol to all-trans retinal and then to the active form all-trans retinoic acid (PubMed:10681376). May also participate in eicosanoids metabolism by converting hydroperoxide species into oxo metabolites (lipoxygenase-like reaction, NADPH-independent) (PubMed:21068195).
Cellular Location	Endoplasmic reticulum membrane {ECO:0000250 UniProtKB:P00185}; Peripheral membrane protein {ECO:0000250 UniProtKB:P00185}. Mitochondrion inner membrane {ECO:0000250 UniProtKB:P00185}; Peripheral membrane protein {ECO:0000250 UniProtKB:P00185}. Microsome membrane {ECO:0000250 UniProtKB:P00185}; Peripheral membrane protein {ECO:0000250 UniProtKB:P00185}. Cytoplasm {ECO:0000250 UniProtKB:P00185}
Tissue Location	Lung, lymphocytes and placenta.

Background


CYP1A1 is involved in phase I xenobiotic and drug metabolism (one substrate of it is theophylline). It is inhibited by fluoroquinolones and macrolides and induced by aromatic hydrocarbons. CYP1A1 is also known as AHH (aryl hydrocarbon hydroxylase). It is involved in the metabolic activation of aromatic hydrocarbons (polycyclic aromatic hydrocarbons, PAH), for example, benzo(a)pyrene (BP), by transforming it to an epoxide.

In this reaction, the oxidation of benzo[a]pyrene is catalysed by CYP1A1 to form BP-7,8-epoxide, which can be further oxidized by epoxide hydrolase (EH) to form BP-7,8-dihydrodiol. Finally CYP1A1 catalyses this intermediate to form BP-7,8-dihydrodiol-9,10-epoxide, which is the ultimate carcinogen. However, an in vivo experiment with gene-deficient mice has found that the hydroxylation of benzo(a)pyrene by CYP1A1 can have an overall protective effect on the DNA, rather than contributing to potentially carcinogenic DNA modifications. This effect is likely due to the fact that CYP1A1 is highly active in the intestinal mucosa, and thus inhibits infiltration of ingested benzo(a)pyrene carcinogen into the systemic circulation.


Images




Anti- CYP1A1 Picoband antibody, ABO12231, Western blotting
All lanes: Anti CYP1A1 (ABO12231) at 0.5ug/ml
Lane 1: Rat Lung Tissue Lysate at 50ug
Lane 2: Mouse Lung Tissue Lysate at 50ug
Lane 3: Human Placenta Tissue Lysate at 50ug
Lane 4: JURKAT Whole Cell Lysate at 40ug
Predicted bind size: 58KD
Observed bind size: 58KD



Anti- CYP1A1 Picoband antibody, ABO12231, IHC(P)
IHC(P): Mouse Kidney Tissue





Anti- CYP1A1 Picoband antibody, ABO12231, IHC(P)
IHC(P): Rat Kidney Tissue



Anti- CYP1A1 Picoband antibody, ABO12231, IHC(P)
IHC(P): Human Mammary Cancer Tissue

Anti- CYP1A1 Picoband antibody, ABO12231, IHC(F)
IHC(F): Mouse Kidney Tissue



Anti- CYP1A1 Picoband antibody, ABO12231,IHC(F)IHC(F):
Human Placenta Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.