

Anti-CCL4/MIP-1 beta Antibody

Catalog # ABO12258

Product Information

Application	WB, E
Primary Accession	P13236
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for C-C motif chemokine 4(CCL4) detection. Tested with WB, ELISA in Human.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	388372;6351
Other Names	C-C motif chemokine 4, G-26 T-lymphocyte-secreted protein, HC21, Lymphocyte activation gene 1 protein, LAG-1, MIP-1-beta(1-69), Macrophage inflammatory protein 1-beta, MIP-1-beta, PAT 744, Protein H400, SIS-gamma, Small-inducible cytokine A4, T-cell activation protein 2, ACT-2, MIP-1-beta(3-69), CCL4, LAG1, MIP1B, SCYA4
Calculated MW	10212
Application Details	ELISA , 0.1-0.5 μ g/ml, Human, - Western blot, 0.1-0.5 μ g/ml, Human
Subcellular Localization	Secreted.
Source	Eukaryota
Protein Name	C-C motif chemokine 4
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E. coli-derived human CCL4 recombinant protein (Position: A24-N92). Human CCL4 shares 76.8% and 79.7% amino acid (aa) sequence identity with mouse and rat CCL4, respectively.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also

Sequence Similarities

be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.

Belongs to the intercrine beta (chemokine CC) family.

Protein Information

Name	CCL4
Synonyms	LAG1, MIP1B, SCYA4
Function	Monokine with inflammatory and chemokinetic properties. Binds to CCR5. One of the major HIV-suppressive factors produced by CD8+ T- cells. Recombinant MIP-1-beta induces a dose-dependent inhibition of different strains of HIV-1, HIV-2, and simian immunodeficiency virus (SIV). The processed form MIP-1-beta(3-69) retains the abilities to induce down-modulation of surface expression of the chemokine receptor CCR5 and to inhibit the CCR5-mediated entry of HIV-1 in T-cells. MIP-1- beta(3-69) is also a ligand for CCR1 and CCR2 isoform B.
Cellular Location	Secreted.

Background

Chemokine(C-C motif) ligand 4, also known as CCL4, is a protein which in human is encoded by the CCL4 gene. By analysis of somatic cell hybrids and by in situ hybridization, the CCL4 gene was assigned to 17q21-q23. It is a CC chemokine with specificity for CCR5 receptors. And it is a chemoattractant for natural killer cells, monocytes and a variety of other immune cells. CCL4 is a major HIV-suppressive factor produced by CD8+ T cells. Performing-low memory CD8+ T cells that normally synthesize CCL4. Based on functional analysis of CCL4, one or more additional beta chemokine receptors that transduce CCL4 signals must also exist. MIP-1alpha and CCL4 have been shown to lack affinity for the erythrocyte surface chemokine receptor/Duffy antigen that binds many alpha as well as beta chemokines(11, 14).

Images



Anti- CCL4 Picoband antibody, ABO12258, Western blottingAll lanes: Anti CCL4 (ABO12258) at 0.5ug/mlWB: HELA Whole Cell Lysate at 40ugPredicted bind size: 11KDObserved bind size: 11KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.