

Anti-G-CSF Picoband Antibody

Catalog # ABO12263

Product Information

Application	WB, E
Primary Accession	P09920
Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Granulocyte colony-stimulating factor(CSF3) detection. Tested with WB, ELISA in Human;Mouse.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	12985
Other Names	Granulocyte colony-stimulating factor, G-CSF, Csf3, Csfg
Calculated MW	22421
Application Details	Western blot, 0.1-0.5 μ g/ml, Human, Mouse ELISA , 0.1-0.5 μ g/ml, Mouse
Subcellular Localization	Secreted.
Source	Eukaryota
Protein Name	Granulocyte colony-stimulating factor
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E. coli-derived mouse G-CSF recombinant protein (Position: R47-A208). Mouse G-CSF shares 77.2% amino acid (aa) sequence identity with human G-CSF.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.

Protein Information

Name	Csf3
Synonyms	Csfg
Function	Granulocyte/macrophage colony-stimulating factors are cytokines that act in hematopoiesis by controlling the production, differentiation, and function of 2 related white cell populations of the blood, the granulocytes and the monocytes-macrophages. This CSF induces granulocytes.
Cellular Location	Secreted.

Background

Granulocyte-colony stimulating factor (G-CSF or GCSF), also known as colony-stimulating factor 3 (CSF 3), is a glycoprotein that stimulates the bone marrow to produce granulocytes and stem cells and release them into the bloodstream. Functionally, it is a cytokine and hormone, a type of colony-stimulating factor, and is produced by a number of different tissues. The pharmaceutical analogs of naturally occurring G-CSF are called filgrastim and lenograstim. G-CSF also stimulates the survival, proliferation, differentiation, and function of neutrophil precursors and mature neutrophils.

Images



Anti- G-CSF Picoband antibody, ABO12263, Western blotting
All lanes: Anti G-CSF (ABO12263) at 0.5ug/ml
WB: HELA Whole Cell Lysate at 40ug
Predicted bind size: 22KD
Observed bind size: 22KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.