

Anti-DLD Picoband Antibody

Catalog # ABO12265

Product Information

Application	WB, IHC-P
Primary Accession	P09622
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Dihydrolipoyl dehydrogenase, mitochondrial(DLD) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	1738
Other Names	Dihydrolipoyl dehydrogenase, mitochondrial, 1.8.1.4, Dihydrolipoamide dehydrogenase, Glycine cleavage system L protein, DLD, GCSL, LAD, PHE3
Calculated MW	54177
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, By Heat Western blot, 0.1-0.5 μ g/ml, Human, Mouse, Rat
Subcellular Localization	Mitochondrion matrix.
Source	Eukaryota
Protein Name	Dihydrolipoyl dehydrogenase, mitochondrial
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E.coli-derived human DLD recombinant protein (Position: K300-F509). Human DLD shares 96.2% and 95.7% amino acid (aa) sequence identity with mouse and rat DLD, respectively.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated

Sequence Similarities

freezing and thawing.

Belongs to the class-I pyridine nucleotide-disulfide oxidoreductase family.

Protein Information

Name DLD ([HGNC:2898](#))

Synonyms GCSL, LAD, PHE3

Function Lipoamide dehydrogenase is a component of the glycine cleavage system as well as an E3 component of three alpha-ketoacid dehydrogenase complexes (pyruvate-, alpha-ketoglutarate-, and branched- chain amino acid-dehydrogenase complex) (PubMed:[15712224](#), PubMed:[16442803](#), PubMed:[16770810](#), PubMed:[17404228](#), PubMed:[20160912](#), PubMed:[20385101](#)). The 2-oxoglutarate dehydrogenase complex is mainly active in the mitochondrion (PubMed:[29211711](#)). A fraction of the 2-oxoglutarate dehydrogenase complex also localizes in the nucleus and is required for lysine succinylation of histones: associates with KAT2A on chromatin and provides succinyl-CoA to histone succinyltransferase KAT2A (PubMed:[29211711](#)). In monomeric form may have additional moonlighting function as serine protease (PubMed:[17404228](#)). Involved in the hyperactivation of spermatazoa during capacitation and in the spermatazoal acrosome reaction (By similarity). The pyruvate dehydrogenase (PDH) complex catalyzes the overall conversion of pyruvate to acetyl-CoA and CO(2), and thereby links cytoplasmic glycolysis and the mitochondrial tricarboxylic acid (TCA) cycle (Probable). It contains multiple copies of three enzymatic components: pyruvate dehydrogenase (E1), dihydrolipoamide acetyltransferase (E2) and dihydrolipoamide dehydrogenase (E3) (Probable). The E3 subunit catalyzes reoxidation of the dihydrolipoyl moiety on lipoyl-bearing domains (LBDs) of E2 with NAD+ as the ultimate electron acceptor (PubMed:[16442803](#), PubMed:[16770810](#), PubMed:[20160912](#), PubMed:[20385101](#)).


Cellular Location Mitochondrion matrix. Nucleus. Cell projection, cilium, flagellum {ECO:0000250|UniProtKB:Q811C4}. Cytoplasmic vesicle, secretory vesicle, acrosome. Note=Mainly localizes in the mitochondrion. A small fraction localizes to the nucleus, where the 2- oxoglutarate dehydrogenase complex is required for histone succinylation.

Background


DLD, Dihydrolipoamide dehydrogenase, is a component of the pyruvate dehydrogenase complex, the alpha-ketoglutarate dehydrogenase complex, and the branched-chain alpha-keto acid dehydrogenase complex (BCKD). DLD is a flavoprotein enzyme that degrades lipoamide, and produces dihydrolipoamide. The DLD gene contains 14 exons. The gene is localized to 7q31-q32. This gene encodes the L protein of the mitochondrial glycine cleavage system. The L protein, also named dihydrolipoamide dehydrogenase, is also a component of the pyruvate dehydrogenase complex, the alpha-ketoglutarate dehydrogenase complex, and the branched-chain alpha-keto acid dehydrogenase complex.

Images

Anti- DLD Picoband antibody, ABO12265, Western blottingAll lanes: Anti DLD (ABO12265) at 0.5ug/mlLane 1: Rat Brain Tissue Lysate at 50ugLane 2: Rat Liver Tissue Lysate at 50ugLane 3: Rat Testis Tissue Lysate at



50ugLane 4: Mouse Ovary Tissue Lysate at 50ugLane 5:
 HELA Whole Cell Lysate at 40ugLane 6: SMMC Whole Cell
 Lysate at 40ugPredicted bind size: 54KDObserved bind
 size: 54KD



Anti- DLD Picoband antibody, ABO12265, IHC(P)IHC(P):
 Human Intestinal Cancer Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.