

Anti-KIN Picoband Antibody

Catalog # ABO12293

Product Information

Application	WB, IHC-P
Primary Accession	O60870
Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for DNA/RNA-binding protein KIN17(KIN) detection. Tested with WB, IHC-P in Human;Mouse.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	22944
Other Names	DNA/RNA-binding protein KIN17, Binding to curved DNA, KIN, antigenic determinant of recA protein homolog, KIN (HGNC:6327), BTCDD, KIN17
Calculated MW	45374
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat Western blot, 0.1-0.5 µg/ml, Human, Mouse
Subcellular Localization	Nucleus . Cytoplasm . During S phase, strongly associated with the nuclear matrix, and to chromosomal DNA in the presence of DNA damage. Also shows cytoplasmic localization in elongated spermatids. .
Tissue Specificity	Ubiquitously expressed in all tissues examined, with highest levels in skeletal muscle, heart and testis. Differentially expressed in non-tumorigenic and tumorigenic cell lines. Highly expressed in proliferating epithelial keratinocyte cells in vitro (at protein level). .
Source	Eukaryota
Protein Name	DNA/RNA-binding protein KIN17
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	E.coli-derived human KIN recombinant protein (Position: K3-K202). Human KIN shares 98% amino acid (aa) sequence identity with mouse KIN.

Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20° C for one year. After r° Constitution, at 4° C for one month. It° Can also be aliquotted and stored frozen at -20° C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the KIN17 family.

Protein Information


Name	KIN (HGNC:6327)
Synonyms	BTCD, KIN17
Function	Involved in DNA replication and the cellular response to DNA damage. May participate in DNA replication factories and create a bridge between DNA replication and repair mediated by high molecular weight complexes. May play a role in illegitimate recombination and regulation of gene expression. May participate in mRNA processing. Binds, in vitro, to double-stranded DNA. Also shown to bind preferentially to curved DNA in vitro and in vivo (By similarity). Binds via its C-terminal domain to RNA in vitro.
Cellular Location	Nucleus. Cytoplasm. Note=During S phase, strongly associated with the nuclear matrix, and to chromosomal DNA in the presence of DNA damage. Also shows cytoplasmic localization in elongated spermatids {ECO:0000250 UniProtKB:Q8K339, ECO:0000269 PubMed:11880372, ECO:0000269 PubMed:12359749, ECO:0000269 PubMed:12754299, ECO:0000269 PubMed:12853634, ECO:0000269 PubMed:15831485}
Tissue Location	Ubiquitously expressed in all tissues examined, with highest levels in skeletal muscle, heart and testis Differentially expressed in non-tumorigenic and tumorigenic cell lines Highly expressed in proliferating epithelial keratinocyte cells in vitro (at protein level).

Background


DNA/RNA-binding protein KIN17, also known as BTCD or KIN17, is a protein that in humans is encoded by the KIN gene. This gene is mapped to 10p14. The protein encoded by this gene is a nuclear protein that forms intranuclear foci during proliferation and is redistributed in the nucleoplasm during the cell cycle. Short-wave ultraviolet light provokes the relocalization of the protein, suggesting its participation in the cellular response to DNA damage. Originally selected based on protein-binding with RecA antibodies, the mouse protein presents a limited similarity with a functional domain of the bacterial RecA protein, a characteristic shared by this human ortholog. Alternative splicing of this gene results in multiple transcript variants.

Images

Anti- KIN Picoband antibody, ABO12293, Western blottingAll lanes: Anti KIN (ABO12293) at 0.5ug/mlLane 1: HELA Whole Cell Lysate at 40ugLane 2: 22RV1 Whole Cell Lysate at 40ugLane 3: COLO320 Whole Cell Lysate at 40ugLane 4: MCF-7 Whole Cell Lysate at 40ugLane 5: U2OS Whole Cell Lysate at 40ugLane 6: NIH3T3 Whole Cell Lysate at 40ugPredicted bind size: 45KDObserved bind



size: 45KD



Anti- KIN Picoband antibody, ABO12293, IHC(P)IHC(P):
Human Oesophagus Squama Cancer Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.