

Anti-HINT1 Picoband Antibody

Catalog # ABO12318

Product Information

Application	WB, IHC-P
Primary Accession	P49773
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Histidine triad nucleotide-binding protein 1(HINT1) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	3094
Other Names	Histidine triad nucleotide-binding protein 1, 3-...-, Adenosine 5'-monophosphoramidase, Protein kinase C inhibitor 1, Protein kinase C-interacting protein 1, PKCI-1, HINT1, HINT, PKCI1, PRKCNH1
Calculated MW	13802
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5 µg/ml, Human, Rat
Subcellular Localization	Cytoplasm. Nucleus. Interaction with CDK7 leads to a more nuclear localization.
Tissue Specificity	Widely expressed.
Source	Eukaryota
Protein Name	Histidine triad nucleotide-binding protein 1
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human HINT1 (59-92aa HISQISVAEDDDDESLLGHLMI VG KKCAADLGLNK), different from the related mouse and rat sequences by three amino acids.
Purification	Immunogen affinity purified.

Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.


Protein Information

Name	HINT1
Synonyms	HINT, PKCI1, PRKCNH1
Function	Exhibits adenosine 5'-monophosphoramidase activity, hydrolyzing purine nucleotide phosphoramidates with a single phosphate group such as adenosine 5'monophosphoramidate (AMP-NH ₂) to yield AMP and NH ₂ (PubMed: 15703176 , PubMed: 16835243 , PubMed: 17217311 , PubMed: 17337452 , PubMed: 22329685 , PubMed: 23614568 , PubMed: 28691797 , PubMed: 29787766 , PubMed: 31990367). Hydrolyzes adenosine 5'monophosphomorpholidate (AMP-morpholidate) and guanosine 5'monophosphomorpholidate (GMP-morpholidate) (PubMed: 15703176 , PubMed: 16835243). Hydrolyzes lysyl-AMP (AMP-N-epsilon-(N-alpha-acetyl lysine methyl ester)) generated by lysine tRNA ligase, as well as Met- AMP, His-AMP and Asp-AMP, lysyl-GMP (GMP-N-epsilon-(N-alpha-acetyl lysine methyl ester)) and AMP-N-alanine methyl ester (PubMed: 15703176 , PubMed: 17337452 , PubMed: 22329685). Hydrolyzes 3-indolepropionic acyl-adenylate, tryptamine adenosine phosphoramidate monoester and other fluorogenic purine nucleoside tryptamine phosphoramidates in vitro (PubMed: 17217311 , PubMed: 17337452 , PubMed: 23614568 , PubMed: 28691797 , PubMed: 29787766 , PubMed: 31990367). Can also convert adenosine 5'-O- phosphorothioate and guanosine 5'-O-phosphorothioate to the corresponding nucleoside 5'-O-phosphates with concomitant release of hydrogen sulfide (PubMed: 30772266). In addition, functions as scaffolding protein that modulates transcriptional activation by the LEF1/TCF1-CTNNB1 complex and by the complex formed with MITF and CTNNB1 (PubMed: 16014379 , PubMed: 22647378). Modulates p53/TP53 levels and p53/TP53-mediated apoptosis (PubMed: 16835243). Modulates proteasomal degradation of target proteins by the SCF (SKP2-CUL1-F-box protein) E3 ubiquitin-protein ligase complex (PubMed: 19112177). Also exhibits SUMO-specific isopeptidase activity, deconjugating SUMO1 from RGS17 (PubMed: 31088288). Deconjugates SUMO1 from RANGAP1 (By similarity).
Cellular Location	Cytoplasm. Nucleus. Note=Interaction with CDK7 leads to a more nuclear localization.
Tissue Location	Widely expressed.


Background

Histidine triad nucleotide-binding protein 1 is a protein that in humans is encoded by the HINT1 gene. It is a haploinsufficient tumor suppressor gene that inhibits the Wnt/ β -catenin pathway in colon cancer cells and Microphthalmia-associated transcription factor (MITF) activity in human mast cells. The protein encoded by this gene can hydrolyze substrates such as AMP-morpholidate, AMP-N-alanine methyl ester, AMP-alpha-acetyl lysine methyl ester, and AMP-NH₂. The encoded protein interacts with these substrates via a histidine triad motif, which is part of the loop that binds to the substrate. Several transcript variants, but only one of them protein-coding, have been found for this gene.


Images




Anti- HINT1 Picoband antibody, ABO12318, Western blotting
 All lanes: Anti HINT1 (ABO12318) at 0.5ug/ml
 Lane 1: Rat Liver Tissue Lysate at 50ug
 Lane 2: Rat Thymus Tissue Lysate at 50ug
 Lane 3: Rat Testis Tissue Lysate at 50ug
 Lane 4: Rat Skeletal Muscle Tissue Lysate at 50ug
 Lane 5: HELA Whole Cell Lysate at 40ug
 Lane 6: 22RV1 Whole Cell Lysate at 40ug
 Predicted bind size: 14KD
 Observed bind size: 14KD



Anti- HINT1 Picoband antibody, ABO12318, IHC(P)
 IHC(P): Mouse Intestine Tissue



Anti- HINT1 Picoband antibody, ABO12318, IHC(P)
 IHC(P): Rat Kidney Tissue



Anti- HINT1 Picoband antibody, ABO12318, IHC(P)
 IHC(P): Human Intestinal Cancer Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.