

Anti-GRP94 Picoband Antibody

Catalog # ABO12323

Product Information

Application	WB
Primary Accession	P14625
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Endoplasmin(HSP90B1) detection. Tested with WB in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	7184
Other Names	Endoplasmin, 94 kDa glucose-regulated protein, GRP-94, Heat shock protein 90 kDa beta member 1, Tumor rejection antigen 1, gp96 homolog, HSP90B1, GRP94, TRA1
Calculated MW	92469
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat
Subcellular Localization	Endoplasmic reticulum lumen. Melanosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.
Source	Eukaryota
Protein Name	Endoplasmin
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	E.coli-derived human GRP94 recombinant protein (Position: R43-H221). Human GRP94 shares 99.4% and 98.9% amino acid (aa) sequence identity with mouse and rat GRP94, respectively.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.


Protein Information

Name	HSP90B1 {ECO:0000303 PubMed:39509507, ECO:0000312 HGNC:HGNC:12028}
Function	ATP-dependent chaperone involved in the processing of proteins in the endoplasmic reticulum, regulating their transport (PubMed: 23572575 , PubMed: 39509507). Together with MESD, acts as a modulator of the Wnt pathway by promoting the folding of LRP6, a coreceptor of the canonical Wnt pathway (PubMed: 23572575 , PubMed: 39509507). When associated with CNPY3, required for proper folding of Toll-like receptors (PubMed: 11584270). Promotes folding and trafficking of TLR4 to the cell surface (PubMed: 11584270). May participate in the unfolding of cytosolic leaderless cargos (lacking the secretion signal sequence) such as the interleukin 1/IL-1 to facilitate their translocation into the ERGIC (endoplasmic reticulum- Golgi intermediate compartment) and secretion; the translocation process is mediated by the cargo receptor TMED10 (PubMed: 32272059).
Cellular Location	Endoplasmic reticulum lumen. Sarcoplasmic reticulum lumen {ECO:0000250 UniProtKB:P41148}. Melanosome Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV.

Background

Heat shock protein 90kDa beta member 1 (HSP90B1), known as endoplasmin, or GRP94, is a chaperone protein that in humans is encoded by the HSP90B1 gene. It is mapped to chromosome 12q23.3. This gene encodes a member of a family of adenosine triphosphate (ATP)-metabolizing molecular chaperones with roles in stabilizing and folding other proteins. The encoded protein is localized to melanosomes and the endoplasmic reticulum. Expression of this protein is associated with a variety of pathogenic states, including tumor formation.

Images



Anti- GRP94 Picoband antibody, ABO12323, Western blotting
All lanes: Anti GRP94 (ABO12323) at 0.5ug/ml
Lane 1: Rat Liver Tissue Lysate at 50ug
Lane 2: A375 Whole Cell Lysate at 40ug
Lane 3: HELA Whole Cell Lysate at 40ug
Lane 4: NIH3T3 Whole Cell Lysate at 40ug
Predicted bind size: 92KD
Observed bind size: 92KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.