

Anti-HSPA2 Picoband Antibody

Catalog # ABO12325

Product Information

Application	WB, IHC-P
Primary Accession	P54652
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Heat shock-related 70 kDa protein 2(HSPA2) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	3306
Other Names	Heat shock-related 70 kDa protein 2, Heat shock 70 kDa protein 2, HSPA2
Calculated MW	70021
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat
Source	Eukaryota
Protein Name	Heat shock-related 70 kDa protein 2
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human HSPA2 (564-598aa KISEQDKNKILDKCQEVINWLDRNQMAEKDEYEHK), identical to the related mouse and rat sequences.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.


Protein Information

Name	HSPA2
Function	Molecular chaperone implicated in a wide variety of cellular processes, including protection of the proteome from stress, folding and transport of newly synthesized polypeptides, activation of proteolysis of misfolded proteins and the formation and dissociation of protein complexes. Plays a pivotal role in the protein quality control system, ensuring the correct folding of proteins, the re-folding of misfolded proteins and controlling the targeting of proteins for subsequent degradation. This is achieved through cycles of ATP binding, ATP hydrolysis and ADP release, mediated by co-chaperones. The affinity for polypeptides is regulated by its nucleotide bound state. In the ATP-bound form, it has a low affinity for substrate proteins. However, upon hydrolysis of the ATP to ADP, it undergoes a conformational change that increases its affinity for substrate proteins. It goes through repeated cycles of ATP hydrolysis and nucleotide exchange, which permits cycles of substrate binding and release (PubMed: 26865365). Plays a role in spermatogenesis. In association with SHCBP1L may participate in the maintenance of spindle integrity during meiosis in male germ cells (By similarity).
Cellular Location	Cytoplasm, cytoskeleton, spindle {ECO:0000250 UniProtKB:P17156}. Note=Colocalizes with SHCBP1L at spindle during the meiosis process. {ECO:0000250 UniProtKB:P17156}

Background


HSPA2 (heat shock 70kDa protein 2) is also known as HEAT-SHOCK PROTEIN, 70-KD, 2, HSP70-2, HEAT-SHOCK PROTEIN, 70-KD, 3 or HSP70-3. Analysis of the sequence indicated that HSPA2 is the human homolog of the murine Hsp70-2 gene, with 91.7% identity in the nucleotide coding sequence and 98.2% in the corresponding amino acid sequence. HSPA2 has less amino acid homology to the other members of the human HSP70 gene family. HSPA2 is constitutively expressed in most tissues, with very high levels in testis and skeletal muscle. The HSPA2 gene is located on chromosome 14q22-q24. Immunohistochemical analysis detected weak expression of HSPA2 in spermatocytes and stronger expression in spermatids and in the tail of mature sperm. HSPA2 may be critical to sperm maturation through its role as a protein chaperone.

Images





Anti- HSPA2 Picoband antibody, ABO12325, Western blotting
All lanes: Anti HSPA2 (ABO12325) at 0.5ug/ml
Lane 1: Rat Liver Tissue Lysate at 50ug
Lane 2: Rat Thymus Tissue Lysate at 50ug
Lane 3: Rat Testis Tissue Lysate at 50ug
Lane 4: Mouse Liver Tissue Lysate at 50ug
Lane 5: Mouse Kidney Tissue Lysate at 50ug
Lane 6: HELA Whole Cell Lysate at 40ug
Lane 7: MCF-7 Whole Cell Lysate at 40ug
Lane 8: A375 Whole Cell Lysate at 40ug
Lane 9: NIH3T3 Whole Cell Lysate at 40ug
Predicted bind size: 70KD
Observed bind size: 70KD

Anti- HSPA2 Picoband antibody, ABO12325, IHC(P)
IHC(P): Mouse Intestine Tissue



Anti- HSPA2 Picoband antibody, ABO12325,IHC(P)IHC(P):
Rat Intestine Tissue



Anti- HSPA2 Picoband antibody, ABO12325,IHC(P)IHC(P):
Human Lung Cancer Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.