

Anti-Galectin 8 Picoband Antibody

Catalog # ABO12345

Product Information

Application	WB
Primary Accession	O00214
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Galectin-8(LGALS8) detection. Tested with WB in Human;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	3964
Other Names	Galectin-8, Gal-8, Po66 carbohydrate-binding protein, Po66-CBP, Prostate carcinoma tumor antigen 1, PCTA-1, LGALS8
Calculated MW	35808
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Rat
Subcellular Localization	Cytoplasm .
Tissue Specificity	Ubiquitous. Selective expression by prostate carcinomas versus normal prostate and benign prostatic hypertrophy.
Source	Eukaryota
Protein Name	Galectin-8
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Galectin 8 (286-317aa HSLEYKHRFKELSSIDTLEINGDIHLLEVRSW), different from the related mouse and rat sequences by six amino acids.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated

freezing and thawing.


Protein Information

Name	LGALS8 (HGNC:6569)
Function	Beta-galactoside-binding lectin that acts as a sensor of membrane damage caused by infection and restricts the proliferation of infecting pathogens by targeting them for autophagy (PubMed: 22246324 , PubMed: 28077878). Detects membrane rupture by binding beta-galactoside ligands located on the luminal side of the endosome membrane; these ligands becoming exposed to the cytoplasm following rupture (PubMed: 22246324 , PubMed: 28077878). Restricts infection by initiating autophagy via interaction with CALCOCO2/NDP52 (PubMed: 22246324 , PubMed: 28077878). Required to restrict infection of bacterial invasion such as S.typhimurium (PubMed: 22246324). Also required to restrict infection of Picornaviridae viruses (PubMed: 28077878). Has a marked preference for 3'-O-sialylated and 3'-O-sulfated glycans (PubMed: 21288902).
Cellular Location	Cytoplasmic vesicle. Cytoplasm, cytosol
Tissue Location	Ubiquitous. Selective expression by prostate carcinomas versus normal prostate and benign prostatic hypertrophy

Background

Galectin-8 is a protein of the galectin family that in humans is encoded by the LGALS8 gene. This gene encodes a member of the galectin family. Galectins are beta-galactoside-binding animal lectins with conserved carbohydrate recognition domains. The galectins have been implicated in many essential functions including development, differentiation, cell-cell adhesion, cell-matrix interaction, growth regulation, apoptosis, and RNA splicing. This gene is widely expressed in tumoral tissues and seems to be involved in integrin-like cell interactions. Alternatively spliced transcript variants encoding different isoforms have been identified.

Images



Anti- Galectin 8 Picoband antibody, ABO12345, Western blottingAll lanes: Anti Galectin 8 (ABO12345) at 0.5ug/mlLane 1: Rat Brain Tissue Lysate at 50ugLane 2: Rat Kidney Tissue Lysate at 50ugLane 3: Human Placenta Tissue Lysate at 50ugLane 4: HELA Whole Cell Lysate at 40ugLane 5: A431 Whole Cell Lysate at 40ugPredicted bind size: 36KDObserved bind size: 50KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.