

Anti-NDRG2 Picoband Antibody

Catalog # ABO12358

Product Information

Application	WB, IHC-P
Primary Accession	Q9UN36
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Protein NDRG2(NDRG2) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	57447
Other Names	Protein NDRG2, N-myc downstream-regulated gene 2 protein, Protein Syld709613, NDRG2, KIAA1248, SYLD
Calculated MW	40798
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Mouse, Rat, Human, By Heat Western blot, 0.1-0.5 μ g/ml, Mouse, Rat, Human
Subcellular Localization	Cytoplasm. Cytoplasm, perinuclear region. Cell projection, growth cone . In neurons, seems to concentrate at axonal growth cone. Perinuclear in neurons (By similarity). .
Tissue Specificity	Highly expressed in brain, heart, skeletal muscle and salivary gland, and moderately in kidney and liver. Expressed in dendritic cells, but not in other blood cells. Expression levels are low in pancreatic and liver cancer tissues; absent in meningioma. Expressed in low-grade gliomas but present at low levels in glioblastoma. Isoform 1 and isoform 2 are present in brain neurons and up-regulated in Alzheimer disease (at protein level). .
Source	Eukaryota
Protein Name	Protein NDRG2
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human

NDRG2 (210-247aa NSELIQKYRNIITHAPNLDNIELYWNSYNNRRDLNFER), different from the related mouse and rat sequences by three amino acids.

Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.


Protein Information

Name	NDRG2
Synonyms	KIAA1248, SYLD
Function	Contributes to the regulation of the Wnt signaling pathway. Down-regulates CTNNB1-mediated transcriptional activation of target genes, such as CCND1, and may thereby act as tumor suppressor. May be involved in dendritic cell and neuron differentiation.
Cellular Location	Cytoplasm. Cytoplasm, perinuclear region. Cell projection, growth cone. Note=In neurons, seems to concentrate at axonal growth cone. Perinuclear in neurons (By similarity).
Tissue Location	Highly expressed in brain, heart, skeletal muscle and salivary gland, and moderately in kidney and liver. Expressed in dendritic cells, but not in other blood cells. Expression levels are low in pancreatic and liver cancer tissues; absent in meningioma Expressed in low-grade gliomas but present at low levels in glioblastoma. Isoform 1 and isoform 2 are present in brain neurons and up-regulated in Alzheimer disease (at protein level)


Background

Protein NDRG2, also known as KIAA1248, is a protein that in humans is encoded by the NDRG2 gene. It is a member of the N-myc downregulated gene family which belongs to the alpha/beta hydrolase superfamily. This gene is mapped to 14q11.2. The protein encoded by this gene is a cytoplasmic protein that may play a role in neurite outgrowth. This gene may be involved in glioblastoma carcinogenesis. It contributes to the regulation of the WNT signaling pathway. Also, this gene may be involved in dendritic cell and neuron differentiation.


Images



Anti- NDRG2 Picoband antibody, ABO12358, Western blottingAll lanes: Anti NDRG2 (ABO12358) at 0.5ug/mlLane 1: Rat Brain Tissue Lysate at 50ugLane 2: Mouse Brain Tissue Lysate at 50ugPredicted bind size: 41KDObserved bind size: 41KD



Anti- NDRG2 Picoband antibody, ABO12358, IHC(P)IHC(P):
Mouse Brain Tissue



Anti- NDRG2 Picoband antibody, ABO12358, IHC(P)IHC(P):
Rat Brain Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.