

Anti-IGFBP-3 Antibody

Catalog # ABO12396

Product Information

Application	WB, IHC-P, E
Primary Accession	P17936
Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Insulin-like growth factor-binding protein 3(IGFBP3) detection. Tested with WB, IHC-P, ELISA in Human;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	3486
Other Names	Insulin-like growth factor-binding protein 3, IBP-3, IGF-binding protein 3, IGFBP-3, IGFBP3, IBP3
Calculated MW	31674
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat ELISA , 0.1-0.5 µg/ml, Human, - Western blot, 0.1-0.5 µg/ml, Human, Rat
Subcellular Localization	Secreted .
Tissue Specificity	Expressed by most tissues. Present in plasma.
Source	Eukaryota
Protein Name	Insulin-like growth factor-binding protein 3
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Na ₃ N.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human IGFBP-3 (214-252aa RREMEDTLNHLKFLNVLSPRGVHIPNCDKKGFYKKKQCR), identical to the related mouse and rat sequences.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins

Storage At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.

Protein Information

Name IGFBP3

Synonyms IBP3

Function Multifunctional protein that plays a critical role in regulating the availability of IGFs such as IGF1 and IGF2 to their receptors and thereby regulates IGF-mediated cellular processes including proliferation, differentiation, and apoptosis in a cell-type specific manner (PubMed:[10874028](#), PubMed:[19556345](#)). Also exhibits IGF- independent antiproliferative and apoptotic effects mediated by its receptor TMEM219/IGFBP-3R (PubMed:[20353938](#)). Inhibits the positive effect of humanin on insulin sensitivity (PubMed:[19623253](#)). Promotes testicular germ cell apoptosis (PubMed:[19952275](#)). Acts via LRP- 1/alpha2M receptor, also known as TGF-beta type V receptor, to mediate cell growth inhibition independent of IGF1 (PubMed:[9252371](#)). Mechanistically, induces serine-specific dephosphorylation of IRS1 or IRS2 upon ligation to its receptor, leading to the inhibitory cascade (PubMed:[15371331](#)). In the nucleus, interacts with transcription factors such as retinoid X receptor-alpha/RXRA to regulate transcriptional signaling and apoptosis (PubMed:[10874028](#)).


Cellular Location Secreted. Nucleus

Tissue Location Expressed by most tissues. Present in plasma.

Background


IGFBP3, Insulin-like growth fator-binding protein 3, is a member of the insulin-like growth factor binding protein (IGFBP) family and encodes a protein with an IGFBP domain and a thyroglobulin type-I domain. IGFBP3 is located on chromosome 7. The protein forms a ternary complex with insulin-like growth factor acid-labile subunit (IGFALS) and either insulin-like growth factor (IGF) I or II. In this form, it circulates in the plasma, prolonging the half-life of IGFs and altering their interaction with cell surface receptors. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.

Images



Anti- IGFBP-3 Picoband antibody, ABO12396, Western blottingAll lanes: Anti IGFBP-3 (ABO12396) at 0.5ug/mlLane 1: Rat Kidney Tissue Lysate at 50ugLane 2: Rat Liver Tissue Lysate at 50ugLane 3: SGC Whole Cell Lysate at 40ugLane 4: 22RV1 Whole Cell Lysate at 40ugPredicted bind size: 31KDObserved bind size: 31KD

Anti- IGFBP-3 Picoband antibody, ABO12396, IHC(P)IHC(P): Human Intestinal Cancer Tissue



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.