10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Cytokeratin 19 Picoband Antibody

Catalog # ABO12401

Product Information

Application	WB, IHC-P
Primary Accession	P08727
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Keratin, type I cytoskeletal 19(KRT19) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	3880
Other Names	Keratin, type I cytoskeletal 19, Cytokeratin-19, CK-19, Keratin-19, K19, KRT19
Calculated MW	44106
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 □g/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5 □g/ml, Human
Tissue Specificity	Expressed in a defined zone of basal keratinocytes in the deep outer root sheath of hair follicles. Also observed in sweat gland and mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, ectocervical epithelium (at protein level). Expressed in epidermal basal cells, in nipple epidermis and a defined region of the hair follicle. Also seen in a subset of vascular wall cells in both the veins and artery of human umbilical cord, and in umbilical cord vascular smooth muscle. Observed in muscle fibers accumulating in the costameres of myoplasm at the sarcolemma in structures that contain dystrophin and spectrin. .
Source	Eukaryota
Protein Name	Keratin, type I cytoskeletal 19
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Cytokeratin 19 (334-372aa

	QLAHIQALISGIEAQLGDVRADSERQNQEYQRLMDIKSR), different from the related mouse and rat sequences by nine amino acids.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20˚C for one year. After r˚Constitution, at 4˚C for one month. It˚Can also be aliquotted and stored frozen at -20˚C for a longer time.Avoid repeated freezing and thawing.

Protein Information


Name	KRT19 (HGNC:6436)
Function	Structural component of intermediate filaments in epithelial cells. Forms intermediate filaments by heterodimerizing with the type II keratin KRT8, providing mechanical integrity to epithelial cells (PubMed:16000376). Integrates into the intermediate filament network of skeletal muscle fibers alongside desmin and KRT18, contributing to force transmission (By similarity). Specifically interacts with the actin-binding domain of dystrophin, helping to link the contractile apparatus to dystrophin at the costameres of striated muscle (PubMed:16000376).
Tissue Location	Expressed in a defined zone of basal keratinocytes in the deep outer root sheath of hair follicles. Also observed in sweat gland and mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, ectocervical epithelium (at protein level). Expressed in epidermal basal cells, in nipple epidermis and a defined region of the hair follicle. Also seen in a subset of vascular wall cells in both the veins and artery of human umbilical cord, and in umbilical cord vascular smooth muscle. Observed in muscle fibers accumulating in the costameres of myoplasm at the sarcolemma in structures that contain dystrophin and spectrin.


Background

Keratin, type I cytoskeletal 19 is a protein that in humans is encoded by the KRT19 gene. The protein encoded by this gene is a member of the keratin family. It is specifically expressed in the periderm, the transiently superficial layer that envelops the developing epidermis. The type I cytokeratins are clustered in a region of chromosome 17q12-q21. Due to its high sensitivity, KRT19 is the most used marker for the RT-PCR-mediated detection of tumor cells disseminated in lymph nodes, peripheral blood, and bone marrow of breast cancer patients. Keratin 19 is often used together with keratin 8 and keratin 18 to differentiate cells of epithelial origin from hematopoietic cells in tests that enumerate circulating tumor cells in blood.


Images

Anti- Cytokeratin 19 Picoband antibody, ABO12401, Western blottingAll lanes: Anti Cytokeratin 19 (ABO12401) at 0.5ug/mlLane 1: Human Placenta Tissue Lysate at 50ugLane 2: MCF-7 Whole Cell Lysate at 40ugPredicted bind size: 44KDObserved bind size: 44KD






Anti- Cytokeratin 19 Picoband antibody, ABO12401, IHC(P)IHC(P): Mouse Intestine Tissue



Anti- Cytokeratin 19 Picoband antibody, ABO12401, IHC(P)IHC(P): Rat Kidney Tissue



Anti- Cytokeratin 19 Picoband antibody, ABO12401, IHC(P)IHC(P): Human Mammary Cancer Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.