

Anti-LIMK1 Picoband Antibody

Catalog # ABO12402

Product Information

Application	WB
Primary Accession	P53667
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for LIM domain kinase 1(LIMK1) detection. Tested with WB in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	3984
Other Names	LIM domain kinase 1, LIMK-1, 2.7.11.1, LIMK1, LIMK
Calculated MW	72585
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat
Subcellular Localization	Cytoplasm. Nucleus. Predominantly found in the cytoplasm.
Tissue Specificity	Highest expression in both adult and fetal nervous system. Detected ubiquitously throughout the different regions of adult brain, with highest levels in the cerebral cortex. Expressed to a lesser extent in heart and skeletal muscle.
Source	Eukaryota
Protein Name	LIM domain kinase 1
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human LIMK1 (599-634aa KLEHWLETLRMHLAGHLPLGPQLEQLDRGFWETYRR), different from the related mouse sequence by three amino acids, and from the related rat sequence by two amino acids.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins

Storage	At -20° C for one year. After r° Constitution, at 4° C for one month. It° Can also be aliquotted and stored frozen at -20° C for a longer time.Avoid repeated freezing and thawing.
----------------	---

Protein Information

Name	LIMK1
Synonyms	LIMK
Function	Serine/threonine-protein kinase that plays an essential role in the regulation of actin filament dynamics. Acts downstream of several Rho family GTPase signal transduction pathways (PubMed: 10436159 , PubMed: 11832213 , PubMed: 12807904 , PubMed: 15660133 , PubMed: 16230460 , PubMed: 18028908 , PubMed: 22328514 , PubMed: 23633677). Activated by upstream kinases including ROCK1, PAK1 and PAK4, which phosphorylate LIMK1 on a threonine residue located in its activation loop (PubMed: 10436159). LIMK1 subsequently phosphorylates and inactivates the actin binding/depolymerizing factors cofilin-1/CFL1, cofilin-2/CFL2 and destrin/DSTN, thereby preventing the cleavage of filamentous actin (F-actin), and stabilizing the actin cytoskeleton (PubMed: 11832213 , PubMed: 15660133 , PubMed: 16230460 , PubMed: 23633677). In this way LIMK1 regulates several actin-dependent biological processes including cell motility, cell cycle progression, and differentiation (PubMed: 11832213 , PubMed: 15660133 , PubMed: 16230460 , PubMed: 23633677). Phosphorylates TPPP on serine residues, thereby promoting microtubule disassembly (PubMed: 18028908). Stimulates axonal outgrowth and may be involved in brain development (PubMed: 18028908).
Cellular Location	Cytoplasm. Nucleus. Cytoplasm, cytoskeleton. Cell projection, lamellipodium {ECO:0000250 UniProtKB:P53668} Note=Predominantly found in the cytoplasm. Localizes in the lamellipodium in a CDC42BPA, CDC42BPB and FAM89B/LRAP25-dependent manner. {ECO:0000250 UniProtKB:P53668}
Tissue Location	Highest expression in both adult and fetal nervous system. Detected ubiquitously throughout the different regions of adult brain, with highest levels in the cerebral cortex. Expressed to a lesser extent in heart and skeletal muscle

Background

LIM domain kinase 1 is an enzyme that in humans is encoded by the LIMK1 gene. There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs, a central PDZ domain, and a C-terminal protein kinase domain. LIMK1 is likely to be a component of an intracellular signaling pathway and may be involved in brain development.

Images

Anti- LIMK1 Picoband antibody, ABO12402, Western blottingAll lanes: Anti LIMK1 (ABO12402) at 0.5ug/mlLane 1: Rat Brain Tissue Lysate at 50ugLane 2: Mouse Gaster Tissue Lysate at 50ugLane 3: HELA Whole Cell Lysate at 40ugLane 4: U87 Whole Cell Lysate at 40ugLane 5: SKOV



Whole Cell Lysate at 40ug Predicted bind size:
72KD Observed bind size: 72KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.