

Anti-SOD3 Picoband Antibody

Catalog # ABO12433

Product Information

Application	WB, IHC-P
Primary Accession	O09164
Host	Rabbit
Reactivity	Mouse
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Extracellular superoxide dismutase [Cu-Zn](SOD3) detection. Tested with WB, IHC-P in Mouse.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	20657
Other Names	Extracellular superoxide dismutase [Cu-Zn], EC-SOD, 1.15.1.1, Sod3
Calculated MW	27392
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Mouse, By Heat Western blot, 0.1-0.5 μ g/ml, Mouse
Subcellular Localization	Secreted, extracellular space.
Source	Eukaryota
Protein Name	Extracellular superoxide dismutase [Cu-Zn]
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of mouse SOD3 (37-75aa EKIDRLDLVEKIGDTHAKVLEIWMELGRRREVDAAEMHA), different from the related rat sequence by nine amino acids.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins.
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.


Protein Information

Name	Sod3
Function	Protect the extracellular space from toxic effect of reactive oxygen intermediates by converting superoxide radicals into hydrogen peroxide and oxygen.
Cellular Location	Secreted, extracellular space. Golgi apparatus, trans-Golgi network


Background

SOD3 (SUPEROXIDE DISMUTASE 3), also called SUPEROXIDE DISMUTASE, EXTRACELLULAR, EC-SOD, and Cu-Zn, is an enzyme that in humans is encoded by the SOD3 gene. This gene encodes a member of the superoxide dismutase (SOD) protein family. SODs are antioxidant enzymes that catalyze the dismutation of two superoxide radicals into hydrogen peroxide and oxygen. Hendrickson et al. (1990) mapped the SOD3 gene to 4pter-q21 by a study of somatic cell hybrids. Stern et al. (2003) narrowed the assignment to 4p15.3-p15.1 by somatic cell and radiation hybrid analysis, linkage mapping, and FISH. The product of this gene is thought to protect the brain, lungs, and other tissues from oxidative stress. The protein is secreted into the extracellular space and forms a glycosylated homotetramer that is anchored to the extracellular matrix (ECM) and cell surfaces through an interaction with heparan sulfate proteoglycan and collagen. A fraction of the protein is cleaved near the C-terminus before secretion to generate circulating tetramers that do not interact with the ECM.

Images



Anti- SOD3 Picoband antibody, ABO12433, Western blotting
All lanes: Anti SOD3 (ABO12433) at 0.5ug/ml
Lane 1: Mouse Lung Tissue Lysate at 50ug
Lane 2: Mouse Kidney Tissue Lysate at 50ug
Lane 3: Mouse Cardiac Muscle Tissue Lysate at 50ug
Predicted bind size: 26KD
Observed bind size: 32KD



Anti- SOD3 Picoband antibody, ABO12433, IHC(P)
IHC(P): Mouse Liver Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.