

Anti-CD19 Picoband Antibody

Catalog # ABO12486

Product Information

Application	WB, IHC-P
Primary Accession	P15391
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for B-lymphocyte antigen CD19(CD19) detection. Tested with WB, IHC-P in Human.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	930
Other Names	B-lymphocyte antigen CD19, B-lymphocyte surface antigen B4, Differentiation antigen CD19, T-cell surface antigen Leu-12, CD19, CD19
Calculated MW	61128
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, By Heat Western blot, 0.1-0.5 μ g/ml, Human
Subcellular Localization	Membrane; Single-pass type I membrane protein.
Source	Eukaryota
Protein Name	B-lymphocyte antigen CD19
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human CD19 (307-337aa LVGILHLQRALVLRKRKRMTDPTRRFFKVT), different from the related mouse sequence by seven amino acids.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins.
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated

freezing and thawing.


Protein Information

Name	CD19
Function	Functions as a coreceptor for the B-cell antigen receptor complex (BCR) on B-lymphocytes (PubMed: 29523808). Decreases the threshold for activation of downstream signaling pathways and for triggering B-cell responses to antigens (PubMed: 1373518 , PubMed: 16672701 , PubMed: 2463100). Activates signaling pathways that lead to the activation of phosphatidylinositol 3-kinase and the mobilization of intracellular Ca(2+) stores (PubMed: 12387743 , PubMed: 16672701 , PubMed: 9317126 , PubMed: 9382888). Is not required for early steps during B cell differentiation in the blood marrow (PubMed: 9317126). Required for normal differentiation of B-1 cells (By similarity). Required for normal B cell differentiation and proliferation in response to antigen challenges (PubMed: 1373518 , PubMed: 2463100). Required for normal levels of serum immunoglobulins, and for production of high-affinity antibodies in response to antigen challenge (PubMed: 12387743 , PubMed: 16672701 , PubMed: 9317126).
Cellular Location	Cell membrane; Single-pass type I membrane protein. Membrane raft {ECO:0000250 UniProtKB:P25918}; Single-pass type I membrane protein {ECO:0000250 UniProtKB:P25918}
Tissue Location	Detected on marginal zone and germinal center B cells in lymph nodes (PubMed: 2463100). Detected on blood B cells (at protein level) (PubMed: 16672701 , PubMed: 2463100)

Background


B-lymphocyte antigen CD19, also known as CD19 (Cluster of Differentiation 19), is a protein that in humans is encoded by the CD19 gene. It is found on the surface of B-cells, a type of white blood cell. Lymphocytes proliferate and differentiate in response to various concentrations of different antigens. The ability of the B cell to respond in a specific, yet sensitive manner to the various antigens is achieved with the use of low-affinity antigen receptors. The CD19 gene encodes a cell surface molecule that assembles with the antigen receptor of B lymphocytes in order to decrease the threshold for antigen receptor-dependent stimulation.

Images



Anti-CD19 Picoband antibody, ABO12486, Western blotting
All lanes: Anti CD19 (ABO12486) at 0.5ug/mL
1: RAJI Whole Cell Lysate at 40ug
Lane 2: A549 Whole Cell Lysate at 40ug
Lane 3: MCF-7 Whole Cell Lysate at 40ug
Lane 4: SW620 Whole Cell Lysate at 40ug
Predicted bind size: 90KD
Observed bind size: 90KD

Anti-CD19 Picoband antibody, ABO12486, IHC(P)
IHC(P): Human Tonsil Tissue



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.