

Anti-Lck Picoband Antibody

Catalog # ABO12494

Product Information

Application	WB, IHC-P
Primary Accession	P06239
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Tyrosine-protein kinase Lck(LCK) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	3932
Other Names	Tyrosine-protein kinase Lck, 2.7.10.2, Leukocyte C-terminal Src kinase, LSK, Lymphocyte cell-specific protein-tyrosine kinase, Protein YT16, Proto-oncogene Lck, T cell-specific protein-tyrosine kinase, p56-LCK, LCK
Calculated MW	58001
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5 µg/ml, Human
Subcellular Localization	Cytoplasm . Cell membrane ; Lipid-anchor ; Cytoplasmic side . Present in lipid rafts in an inactive form.
Tissue Specificity	Expressed specifically in lymphoid cells.
Source	Eukaryota
Protein Name	Tyrosine-protein kinase Lck
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Lck (468-506aa ELYQLMRLCWKERPEDRPTFDYLRSLVEDFFTATEGQYQ), different from the related mouse and rat sequences by three amino acids.
Purification	Immunogen affinity purified.

Cross Reactivity	No cross reactivity with other proteins.
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.

Protein Information


Name	LCK
Function	Non-receptor tyrosine-protein kinase that plays an essential role in the selection and maturation of developing T-cells in the thymus and in the function of mature T-cells (PubMed: 2470098). Plays a key role in T-cell antigen receptor (TCR)-linked signal transduction pathways (PubMed: 2470098). Constitutively associated with the cytoplasmic portions of the CD4 and CD8 surface receptors (PubMed: 2470098). Association of the TCR with a peptide antigen-bound MHC complex facilitates the interaction of CD4 and CD8 with MHC class II and class I molecules, respectively, thereby recruiting the associated LCK protein to the vicinity of the TCR-CD3 complex (PubMed: 2470098). LCK then phosphorylates tyrosine residues within the immunoreceptor tyrosine-based activation motifs (ITAM) of the cytoplasmic tails of the TCR-gamma chains and CD3 subunits, initiating the TCR-CD3 signaling pathway (PubMed: 2470098 , PubMed: 40592325). Once stimulated, the TCR recruits the tyrosine kinase ZAP70, that becomes phosphorylated and activated by LCK. Following this, a large number of signaling molecules are recruited, ultimately leading to lymphokine production. LCK also contributes to signaling by other receptor molecules. Associates directly with the cytoplasmic tail of CD2, which leads to hyperphosphorylation and activation of LCK. Also plays a role in the IL2 receptor-linked signaling pathway that controls the T-cell proliferative response. Binding of IL2 to its receptor results in increased activity of LCK. Is expressed at all stages of thymocyte development and is required for the regulation of maturation events that are governed by both pre-TCR and mature alpha beta TCR. Phosphorylates other substrates including RUNX3, PTK2B/PYK2, the microtubule-associated protein MAPT, RHOH or TYROBP. Interacts with FYB2 (PubMed: 27335501).
Cellular Location	Cell membrane; Lipid-anchor; Cytoplasmic side Cytoplasm, cytosol. Note=Present in lipid rafts in an inactive form.
Tissue Location	Expressed specifically in lymphoid cells.

Background


Lck (or lymphocyte-specific protein tyrosine kinase) is a 56 kDa protein that is found inside specialized cells of the immune system called lymphocytes. The human LCK gene is mapped to chromosome 1p35-p32. This gene is a member of the Src family of protein tyrosine kinases (PTKs). The encoded protein is a key signaling molecule in the selection and maturation of developing T-cells. It contains N-terminal sites for myristylation and palmitoylation, a PTK domain, and SH2 and SH3 domains which are involved in mediating protein-protein interactions with phosphotyrosine-containing and proline-rich motifs, respectively. The protein localizes to the plasma membrane and pericentrosomal vesicles, and binds to cell surface receptors, including CD4 and CD8, and other signaling molecules. Multiple alternatively spliced variants, encoding the same protein, have been described.

Images


Anti- Lck Picoband antibody, ABO12494, Western




blottingAll lanes: Anti Lck (ABO12494) at 0.5ug/mlLane 1: HUT Whole Cell Lysate at 40ugLane 2: JURKAT Whole Cell Lysate at 40ugLane 3: RAJI Whole Cell Lysate at 40ugLane 4: CEM Whole Cell Lysate at 40ugLane 5: K562 Whole Cell Lysate at 40ugPredicted bind size: 58KDObserved bind size: 58KD



Anti- Lck Picoband antibody, ABO12494, IHC(P)IHC(P): Mouse Lymphaden Tissue



Anti- Lck Picoband antibody, ABO12494, IHC(P)IHC(P): Rat Lymphaden Tissue



Anti- Lck Picoband antibody, ABO12494, IHC(P)IHC(P): Human Tonsil Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.